

Hardy spaces $H_L^p(\mathbb{R}^n)$ associated with operators
satisfying k -Davies-Gaffney estimates

CAO Jun & YANG DaChun*

*School of Mathematical Sciences, Beijing Normal University, Laboratory of Mathematics
and Complex Systems, Ministry of Education, Beijing 100875, People's Republic of China
Email: caojun1860@mail.bnu.edu.cn, dcyang@bnu.edu.cn*

Received October 17, 2011; accepted February 20, 2012; published online March 22, 2012

Abstract Let L be a one-to-one operator of type ω having a bounded H_∞ functional calculus and satisfying the k -Davies-Gaffney estimates with $k \in \mathbb{N}$. In this paper, the authors introduce the Hardy space $H_L^p(\mathbb{R}^n)$ with $p \in (0, 1]$ associated with L in terms of square functions defined via $\{e^{-t^{2k}L}\}_{t>0}$ and establish their molecular and generalized square function characterizations. Typical examples of such operators include the $2k$ -order divergence form homogeneous elliptic operator L_1 with complex bounded measurable coefficients and the $2k$ -order Schrödinger type operator $L_2 := (-\Delta)^k + V^k$, where Δ is the Laplacian and $0 \leq V \in L_{\text{loc}}^k(\mathbb{R}^n)$. Moreover, as an application, for $i \in \{1, 2\}$, the authors prove that the associated Riesz transform $\nabla^k(L_i^{-1/2})$ is bounded from $H_{L_i}^p(\mathbb{R}^n)$ to $H^p(\mathbb{R}^n)$ for $p \in (n/(n+k), 1]$ and establish the Riesz transform characterizations of $H_{L_1}^p(\mathbb{R}^n)$ for $p \in (rn/(n+kr), 1]$ if $\{e^{-tL_1}\}_{t>0}$ satisfies the $L^r - L^2$ k -off-diagonal estimates with $r \in (1, 2]$. These results when $k := 1$ and $L := L_1$ are known.

Keywords Hardy space, Hardy-Sobolev space, k -Davies-Gaffney estimate, Schrödinger type operator, higher order elliptic operator, semigroup, square function, higher order Riesz transform, molecule

MSC(2010) 42B35, 42B30, 42B25, 35J10, 42B37

Citation Cao J, Yang D C. Hardy spaces $H_L^p(\mathbb{R}^n)$ associated with operators satisfying k -Davies-Gaffney estimates. Sci China Math, 2012, 55, doi: 10.1007/s11425-000-0000-0

1 Introduction

The Hardy spaces, as a suitable substitute of Lebesgue spaces $L^p(\mathbb{R}^n)$, play an important role in various fields of analysis and partial differential equations. It is well known that the Hardy spaces $H^p(\mathbb{R}^n)$ are essentially related to the Laplacian operator $\Delta := \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2}$, which have been intensively studied; see, for example, [18, 26, 27, 48–50] and the references therein.

In recent years, the study of Hardy spaces associated with different differential operators inspires great interests; see, for example, [4, 7, 8, 12–15, 21–25, 29–32, 35, 37–39, 45, 55] and their references. In particular, in [4], when the operator L satisfies a pointwise Poisson upper bound, Auscher, McIntosh and Duong introduced the Hardy space $H_L^1(\mathbb{R}^n)$ associated with L in terms of area integral functions. Later, in [22, 23], Duong and Yan introduced the BMO-type space $\text{BMO}_L(\mathbb{R}^n)$ associated with such an L and proved the dual space of $H_L^1(\mathbb{R}^n)$ is $\text{BMO}_{L^*}(\mathbb{R}^n)$, where L^* denotes the adjoint operator of L in $L^2(\mathbb{R}^n)$. Yan [53] further generalized these results to the Hardy space $H_L^p(\mathbb{R}^n)$ with $p \in (0, 1]$ close to

*Corresponding author

1 and its dual space. Also, the Orlicz-Hardy space and its dual space associated with such an L were studied in [42, 45].

Auscher and Russ [8] studied the Hardy space H_L^1 on strongly Lipschitz domains associated with a second order divergence form elliptic operator L whose heat kernels have the Gaussian upper bounds and regularity. Very recently, Auscher, McIntosh and Russ [7] treated the Hardy space H^1 associated with the Hodge Laplacian on a Riemannian manifold with doubling measure; Hofmann-Mayboroda in [37, 38] and Hofmann-Mayboroda-McIntosh in [39] introduced the Hardy and Sobolev spaces associated with a second order divergence form elliptic operator L on \mathbb{R}^n with bounded measurable complex coefficients and these operators may not have the pointwise heat kernel bounds, while a theory of the Orlicz-Hardy space and its dual space associated with L was independently developed in [43, 44].

Moreover, a theory of Hardy spaces associated with the Schrödinger operators $-\Delta + V$ was well developed, where the nonnegative potential V satisfies the reverse Hölder inequality (see, for example, Dziubański and Zienkiewicz [24, 25] and Yang and Zhou [56] and their references). More generally, for nonnegative self-adjoint operators L satisfying the Davies-Gaffney estimates, Hofmann et al. [35] introduced a new Hardy space $H_L^1(\mathbb{R}^n)$, which was extended to the Orlicz-Hardy space by Jiang and Yang [41]. Recently, the Hardy space $H_{(-\Delta)^2+V^2}^1(\mathbb{R}^n)$ associated with the Schrödinger type operators $(-\Delta)^2 + V^2$ was also studied in [12].

From now on, in what follows of this paper, we *always* let L be a one-to-one operator of type ω having a bounded H_∞ functional calculus and satisfying the k -Davies-Gaffney estimates with $k \in \mathbb{N}$ (see (2.6) below). Motivated by [35, 39], in this paper, we introduce the Hardy space $H_L^p(\mathbb{R}^n)$ with $p \in (0, 1]$ associated with L in terms of the square function defined via $\{e^{-t^{2k}L}\}_{t>0}$ (see (4.1) below) and establish their molecular and generalized square function characterizations. Typical examples of such operators include the $2k$ -order divergence form homogeneous elliptic operator L_1 with complex bounded measurable coefficients and the $2k$ -order Schrödinger type operator $L_2 := (-\Delta)^k + V^k$, where Δ is the Laplacian and $0 \leq V \in L_{\text{loc}}^k(\mathbb{R}^n)$. Moreover, as an application, for $i \in \{1, 2\}$, we prove that the associated Riesz transform $\nabla^k(L_i^{-1/2})$ is bounded from $H_{L_i}^p(\mathbb{R}^n)$ to $H^p(\mathbb{R}^n)$ for $p \in (n/(n+k), 1]$ and establish the Riesz transform characterizations of $H_{L_1}^p(\mathbb{R}^n)$ for $p \in (rn/(n+kr), 1]$ if $\{e^{-tL_1}\}_{t>0}$ satisfies the $L^r - L^2$ k -off-diagonal estimates with $r \in (1, 2]$ (see Definition 6.3 below for the definition). These results when $k := 1$ and $L := L_1$ were already obtained recently by Hofmann-Mayboroda [37, 38], Jiang-Yang [41, 43], and Hofmann-Mayboroda-McIntosh [39].

A new ingredient appearing in this paper is the introduction of the k -Davies-Gaffney estimates with $k \in \mathbb{N}$, which is naturally satisfied by $2k$ -order Schrödinger operators $(-\Delta)^k + V^k$. Via the perturbation technique (see, for example, [10, 11]) and some ideas from the proof of [19, Lemma 2], and using the elliptic condition, we further show that the semigroup $\{e^{-tL_1}\}_{t>0}$ also satisfies the k -Davies-Gaffney estimates.

Another new observation of this paper is that the nonnegative self-adjoint property of operators in [35, 41] can be weakened into the assumption that L has a bounded H_∞ functional calculus. We point out that when this manuscript was in preparation, we learned from Anh and Li [1] that this was also observed by Duong and Li [20].

This paper is organized as follows. In Section 2, we first recall some results on the H_∞ functional calculus and describe some assumptions on operators considered in this paper. In particular, we introduce the notion of k -Davies-Gaffney estimates with $k \in \mathbb{N}$ in (2.6) below. Some examples satisfying these assumptions are also given in this section.

Let L be an operator satisfying assumptions in Section 2. In Section 3, using some ideas from [35, 37–39], we establish some off-diagonal estimates for some families of operators related to L . More precisely, we show that if $\{e^{-tL}\}_{t>0}$ satisfies the k -Davies-Gaffney estimates, then the family $\{(zL)^m e^{-zL}\}_{z \in S_{\ell(\pi/2-\omega)}^0}$ of operators for any $m \in \mathbb{N} \cup \{0\}$ also satisfies the k -Davies-Gaffney estimates in z (see Lemma 3.1), the k -Davies-Gaffney estimates are stable under compositions (see Lemma 3.2) and the family $\{\psi(tL)f(L)\}_{t>0}$ of operators satisfies the k -Davies-Gaffney estimates of order σ (see (3.7) below for the definition), where ψ belongs to the decaying function class $\Psi_{\sigma,\tau}(S_\mu^0)$ as in (2.2) below (see Lemma 3.3 below). Let L_1 be

the $2k$ -order divergence form homogeneous elliptic operator with complex bounded measurable coefficients and L_2 the $2k$ -order Schrödinger type operator. In this section, we also prove that the semigroup $\{e^{-tL_1}\}_{t>0}$ and the family $\{\sqrt{t}\nabla^k e^{-tL_i}\}_{t>0}$ of operators for $i \in \{1, 2\}$ satisfy the k -Davies-Gaffney estimates, respectively, in Propositions 3.4 and 3.6.

In Definition 4.1 of Section 4, we first introduce the Hardy space $H_L^p(\mathbb{R}^n)$ for $p \in (0, 1]$ in terms of the square function S_L defined via $\{e^{-t^{2k}L}\}_{t>0}$ and, in Definition 4.2, the molecular Hardy space $H_{L, \text{mol}, M}^p(\mathbb{R}^n)$ with $M \in (n(1/p - 1/2)/(2k), \infty)$. Then, by using Lemma 3.1, we prove that for each (H_L^p, ϵ, M) -molecule m , $\|S_L(m)\|_{L^p(\mathbb{R}^n)}$ is uniformly bounded (see (4.4) below), which together with a boundedness criteria from [39] (see also Lemma 4.1 below) implies that $H_{L, \text{mol}, M}^p(\mathbb{R}^n) \subset H_L^p(\mathbb{R}^n)$. On the other hand, using the atomic decomposition of the tent space $T^p(\mathbb{R}_+^{n+1})$ and the k -Davies-Gaffney estimate, we obtain that the operator $\pi_{M, L}$ in (4.11) maps any $T^p(\mathbb{R}_+^{n+1})$ -atom into an (H_L^p, ϵ, M) -molecule up to a harmless positive constant multiple in Lemma 4.8 below. Then, by a Calderón reproducing formula, we establish a molecular decomposition of $H_L^p(\mathbb{R}^n)$ which yields another inclusion $H_L^p(\mathbb{R}^n) \subset H_{L, \text{mol}, M}^p(\mathbb{R}^n)$. Thus, we obtain the molecular characterization of $H_L^p(\mathbb{R}^n)$ in Theorem 4.5 below.

Section 5 is devoted to the generalized square function characterization of $H_L^p(\mathbb{R}^n)$. Motivated by [39], we first introduce the generalized square function Hardy space $H_{\psi, L}^p(\mathbb{R}^n)$ for $p \in (0, 1]$ and some $\psi \in \Psi_{\sigma, \tau}(S_\mu^0)$ in Definition 5.1 below. Then, for any $\psi \in \Psi_{\sigma, \tau}(S_\mu^0)$ and all $f \in H_\infty(S_\mu^0)$ (see (2.1) for the definition), we introduce the operators $Q_{\psi, L}$, $\pi_{\psi, L}$ and their composition Q^f (see (5.1), (5.4) and (5.5) for their definitions). Using the k -Davies-Gaffney estimates of order σ for $\{\psi(tL)f(L)\}_{t>0}$ in Lemma 3.3 below, we prove that the operator Q^f is bounded on the tent space $T^p(\mathbb{R}_+^{n+1})$ (see Lemma 5.5), $Q_{\psi, L}$ is bounded from $H_L^p(\mathbb{R}^n)$ to $T^p(\mathbb{R}_+^{n+1})$ and $\pi_{\psi, L}$ is bounded from $T^p(\mathbb{R}_+^{n+1})$ to $H_L^p(\mathbb{R}^n)$ for some ψ (see Lemma 5.6 below). Combining these boundedness and using a Calderón reproducing formula in (5.14), we then obtain the generalized square function characterization of $H_L^p(\mathbb{R}^n)$ in Theorem 5.2, which is used in obtaining the Riesz transform characterization of $H_{L_1}^p(\mathbb{R}^n)$ in Section 6. For all $\alpha \in (0, \infty)$, let L^α be the fractional power with exponent α of L and the Hardy space $H_{L^\alpha}^p(\mathbb{R}^n)$ be defined as in (5.3) below via the square function S_{L^α} as in (5.2). As another application of Theorem 5.2, we then obtain in Corollary 5.3 that $H_{L^\alpha}^p(\mathbb{R}^n) = H_L^p(\mathbb{R}^n)$ with equivalent norms, in particular, $H_{(-\Delta)^k}^p(\mathbb{R}^n) = H^p(\mathbb{R}^n)$ with equivalent norms for all $k \in \mathbb{N}$, where $H^p(\mathbb{R}^n)$ is the classical Hardy space in [26, 49].

Finally, in Section 6, we concentrate on the behavior of the Riesz transforms $\nabla^k L_i^{-1/2}$ on $H_{L_i}^p(\mathbb{R}^n)$ for $i \in \{1, 2\}$. By the gradient estimates of the semigroup $\{e^{-tL_i}\}_{t>0}$ in Proposition 3.6 and the composition rule of k -Davies-Gaffney estimates in Lemma 3.2, we first show that the two families of operators, $\{\nabla^k L_i^{-1/2}(I - e^{-tL_i})^M\}_{t>0}$ and $\{\nabla^k L_i^{-1/2}(tL_i e^{-tL_i})^M\}_{t>0}$ for all $M \in \mathbb{N}$, satisfy some estimates similar to the k -Davies-Gaffney estimates of order M (see Lemma 6.1 below). Then, using these estimates, we prove that for each $(H_{L_i}^p, \epsilon, M)$ -molecule m with $p \in (n/(n+k), 1]$ and $M \in (n(1/p - 1/2)/(2k), \infty)$, $\nabla^k(L_i^{-1/2})(m)$ is a classical $H^p(\mathbb{R}^n)$ -molecule up to a harmless constant multiple, which further implies that Riesz transforms $\nabla^k(L_i^{-1/2})$ are bounded from $H_{L_i}^p(\mathbb{R}^n)$ to the classical Hardy space $H^p(\mathbb{R}^n)$ in Theorem 6.2 below. In the remainder of this section, motivated by [39], by assuming that the semigroup $\{e^{-tL_1}\}_{t>0}$ satisfies the $L^r - L^2$ k -off-diagonal estimates for $r \in (1, 2]$, we then establish the Riesz transform characterization of $H_{L_1}^p(\mathbb{R}^n)$. To this end, we first show in Lemma 6.5 below that $\{tL_1 e^{-tL_1}\}_{t>0}$ also satisfies the $L^r - L^2$ k -off-diagonal estimates. We then recall some known results concerning the homogeneous Triebel-Lizorkin space $\dot{F}_{p, q}^\alpha(\mathbb{R}^n)$ and their atomic characterizations from [16, 34, 51] and [52, Proposition 4.3]. Let $\dot{W}^{k, 2}(\mathbb{R}^n)$ be the homogenous Sobolev space of order k . With the help of these results, we show that if $f \in \dot{W}^{k, 2}(\mathbb{R}^n) \cap \dot{H}^{k, p}(\mathbb{R}^n)$ when $p \in (0, 1]$, then its atomic decomposition converges in both $\dot{W}^{k, 2}(\mathbb{R}^n)$ and $\dot{H}^{k, p}(\mathbb{R}^n)$ (see Lemma 6.9 below). Moreover, by the $L^r - L^2$ k -off-diagonal estimates for $\{tL_1 e^{-tL_1}\}_{t>0}$, we prove that for each $H^{k, p}(\mathbb{R}^n)$ -atom b , $S_1 \sqrt{L_1}(b)$ is uniformly bounded on $L^p(\mathbb{R}^n)$ (see (6.12) below), which, together with the generalized square function characterization of $H_{L_1}^p(\mathbb{R}^n)$ in Theorem 5.2 and Lemma 6.9, shows that $S_1 \sqrt{L_1}$ is bounded from the Hardy-Sobolev space $\dot{H}^{k, p}(\mathbb{R}^n)$ to $L^p(\mathbb{R}^n)$. This, combined with the boundedness of Riesz transforms on $H_{L_1}^p(\mathbb{R}^n)$ in Theorem 6.2, yields

the Riesz transform characterization of $H_{L_1}^p(\mathbb{R}^n)$ in Theorem 6.10 below. We point out that in the proof of the estimate (6.12), we use the embedding result (6.14) below on the homogeneous Triebel-Lizorkin space from [51] and another key fact from [6, Theorem 1.1] that $\|\sqrt{L_1}f\|_{L^2(\mathbb{R}^n)} \lesssim \|\nabla^k f\|_{L^2(\mathbb{R}^n)}$. The latter fact may not be true for L_2 ; see Remark 6.11 below. Thus, it seems that one needs some new ideas to obtain the Riesz characterization of $H_{L_2}^p(\mathbb{R}^n)$.

We now make some conventions on the notation. Throughout the whole paper, we always let $\mathbb{N} := \{1, 2, \dots\}$ and $\mathbb{Z}_+ := \mathbb{N} \cup \{0\}$. Denote the *differential operator* $\frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}}$ simply by ∂^α , where $\alpha := (\alpha_1, \dots, \alpha_n)$ and $|\alpha| := \alpha_1 + \dots + \alpha_n$. We also denote the *2k-order divergence form homogenous elliptic operator with complex bounded measurable coefficients* $(-1)^k \sum_{|\alpha|=|\beta|=k} \partial^\alpha (a_{\alpha,\beta} \partial^\beta)$ by L_1 and the *2k-order Schrödinger type operator* $(-\Delta)^k + V^k$ by L_2 . We use C to denote a *positive constant*, that is independent of the main parameters involved but whose value may differ from line to line, and $C(\alpha, \dots)$ to denote a *positive constant* depending on the parameters α, \dots . *Constants with subscripts*, such as C_0 , do not change in different occurrences. If $f \leq Cg$, we then write $f \lesssim g$; and if $f \lesssim g \lesssim f$, we then write $f \sim g$. For all $x \in \mathbb{R}^n$ and $r \in (0, \infty)$, let $B(x, r) := \{y \in \mathbb{R}^n : |x - y| < r\}$. Also, for any set $E \in \mathbb{R}^n$, we use E^c to denote $\mathbb{R}^n \setminus E$ and χ_E its *characteristic function*.

2 Preliminaries

We first collect some basic results on the theory of H_∞ functional calculus, developed by McIntosh in [46], that we need in what follows. For more details and further references about functional calculus, we refer the reader to [2, 33, 46] and the references therein.

For $\theta \in [0, \pi)$, the *open and closed sectors*, S_θ^0 and S_θ , of angle θ in the complex plane \mathbb{C} are defined as follows:

$$S_\theta^0 := \{z \in \mathbb{C} \setminus \{0\} : |\arg z| < \theta\}$$

and

$$S_\theta := \{z \in \mathbb{C} \setminus \{0\} : |\arg z| \leq \theta\} \cup \{0\}.$$

Let $\omega \in [0, \pi)$. A closed operator T in $L^2(\mathbb{R}^n)$ is called of *type* ω , if the spectrum of T , $\sigma(T)$, is contained in S_ω , and for each $\theta \in (\omega, \pi)$, there exists a nonnegative constant C such that for all $z \in \mathbb{C} \setminus S_\theta$, $\|(T - zI)^{-1}\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq C|z|^{-1}$, here and in what follows, $\|S\|_{\mathcal{L}(\mathcal{H})}$ denotes the *operator norm* of the linear operator S on the normed linear space \mathcal{H} .

For $\mu \in [0, \pi)$ and $\sigma, \tau \in (0, \infty)$, we need the following spaces of functions:

$$H(S_\mu^0) := \{f : f \text{ is holomorphic on } S_\mu^0\},$$

$$H_\infty(S_\mu^0) := \left\{f \in H(S_\mu^0) : \|f\|_{L^\infty(S_\mu^0)} < \infty\right\} \quad (2.1)$$

and

$$\Psi_{\sigma,\tau}(S_\mu^0) := \{f \in H(S_\mu^0) : |f(\xi)| \leq C \inf\{|\xi|^\sigma, |\xi|^{-\tau}\} \text{ for all } \xi \in S_\mu^0\}. \quad (2.2)$$

It is known that every one-to-one operator T of type ω in $L^2(\mathbb{R}^n)$ has a unique holomorphic functional calculus which is consistent with the usual definition of polynomials of operators (see, for example, [46]). More precisely, let T be a one-to-one operator of type ω , with $\omega \in [0, \pi)$, $\mu \in (\omega, \pi)$, $\sigma, \tau \in (0, \infty)$, and $f \in \Psi_{\sigma,\tau}(S_\mu^0)$. The function of the operator T , $f(T)$ can be defined by the H_∞ functional calculus in the following way,

$$f(T) := \frac{1}{2\pi i} \int_\gamma (\xi I - T)^{-1} f(\xi) d\xi, \quad (2.3)$$

where $\gamma := \{re^{i\nu} : \infty > r > 0\} \cup \{re^{-i\nu} : 0 < r < \infty\}$, $\nu \in (\omega, \mu)$, is a curve consisting of two rays parameterized anti-clockwise. It is well known that the above definition is independent of the choice of $\nu \in (\omega, \mu)$ and the integral in (2.3) is absolutely convergence in $\mathcal{L}(L^2(\mathbb{R}^n))$ (see [33, 46]).

In what follows, we *always assume* $\omega \in [0, \pi/2)$. Then, it follows from [33, Proposition 7.1.1] that for every operator T of type ω in $L^2(\mathbb{R}^n)$, $-T$ generates a holomorphic C_0 -semigroup $\{e^{-zL}\}_{z \in S_{\pi/2-\omega}^0}$ on the open sector $S_{\pi/2-\omega}^0$ such that $\|e^{-zL}\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq 1$ for all $z \in S_{\pi/2-\omega}^0$ and, moreover, every nonnegative self-adjoint operator is of type 0.

Let $\Psi(S_\mu^0) := \cup_{\sigma, \tau > 0} \Psi_{\sigma, \tau}(S_\mu^0)$. By the relationship between the associated semigroup and the resolvent of T , for all $f \in \Psi(S_\mu^0)$, $f(T)$ can further be represented as

$$f(T) = \int_{\Gamma_+} e^{-zT} \eta_+(z) dz + \int_{\Gamma_-} e^{-zT} \eta_-(z) dz, \quad (2.4)$$

where

$$\eta_\pm(z) := \frac{1}{2\pi i} \int_{\gamma_\pm} e^{\xi z} f(\xi) d\xi, \quad z \in \Gamma_\pm, \quad (2.5)$$

$\Gamma_\pm := \mathbb{R}^+ e^{\pm i(\pi/2-\theta)}$, $\gamma_\pm := \mathbb{R}^+ e^{\pm i\nu}$ and $0 \leq \omega < \theta < \nu < \mu < \pi/2$. Here and in what follows, $\mathbb{R}^+ := (0, \infty)$.

It is well known that the above holomorphic functional calculus defined on $\Psi(S_\mu^0)$ can be extended to $H_\infty(S_\mu^0)$ via a limit process (see [46]). Recall that for $\mu \in (0, \pi)$, the operator T is said to *have a bounded $H_\infty(S_\mu^0)$ functional calculus* in the Hilbert space \mathcal{H} , if there exists a positive constant C such that for all $\psi \in H_\infty(S_\mu^0)$, $\|\psi(T)\|_{\mathcal{L}(\mathcal{H})} \leq C\|\psi\|_{L^\infty(S_\mu^0)}$ and T is called to *have a bounded H_∞ functional calculus* in the Hilbert space \mathcal{H} if there exists $\mu \in (0, \pi)$ such that T has a bounded $H_\infty(S_\mu^0)$ functional calculus.

Now, we describe our assumptions of operators L considered in this paper. Throughout the whole paper, we *always assume* that L satisfies the following *assumptions*:

- (A₁) The operator L is a one-to-one operator of type ω in $L^2(\mathbb{R}^n)$ with $\omega \in [0, \pi/2)$;
- (A₂) The operator L has a bounded H_∞ functional calculus in $L^2(\mathbb{R}^n)$;
- (A₃) Let $k \in \mathbb{N}$. The operator L generates a holomorphic semigroup $\{e^{-tL}\}_{t>0}$ which satisfies the *k-Davies-Gaffney estimate*, namely, there exist positive constants \tilde{C} and C_1 such that for all closed sets E and F in \mathbb{R}^n , $t \in (0, \infty)$ and $f \in L^2(\mathbb{R}^n)$ supported in E ,

$$\|e^{-tL}f\|_{L^2(F)} \leq \tilde{C} \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_1 t^{1/(2k-1)}} \right\} \|f\|_{L^2(E)}, \quad (2.6)$$

here and in what follows, $\text{dist}(E, F) := \inf_{x \in E, y \in F} |x - y|$ is the distance between E and F .

Remark 2.1. We point out that when $k = 1$, the *k-Davies-Gaffney estimate* is usually called the *Davies-Gaffney estimate* (or the *L^2 off-diagonal estimate* or just the *Gaffney estimate*) (see, for example, [35, 37–39, 41]).

Let $k \in \mathbb{N}$. Examples of operators, satisfying the above assumptions (A₁), (A₂) and (A₃), include the following $2k$ -order divergence form homogeneous elliptic operator:

$$L_1 := (-1)^k \sum_{|\alpha|=|\beta|=k} \partial^\alpha (a_{\alpha, \beta} \partial^\beta)$$

with complex bounded measurable coefficients $a_{\alpha, \beta}$ for all multi-indices α, β and the $2k$ -order Schrödinger type operator $L_2 := (-\Delta)^k + V^k$ with $0 \leq V \in L_{\text{loc}}^k(\mathbb{R}^n)$. More precisely, let $W^{k,2}(\mathbb{R}^n)$ be the *Sobolev space of order k* endowed with the *norm*

$$\|\cdot\|_{W^{k,2}(\mathbb{R}^n)} := \sum_{0 \leq |\alpha| \leq k} \|\partial^\alpha(\cdot)\|_{L^2(\mathbb{R}^n)}.$$

Denote by \mathfrak{a} the *sesquilinear form* given by

$$\mathfrak{a}(f, g) := \int_{\mathbb{R}^n} \sum_{|\alpha|=|\beta|=k} a_{\alpha,\beta}(x) \partial^\beta f(x) \overline{\partial^\alpha g(x)} dx \quad (2.7)$$

with domain $D(\mathfrak{a}) := W^{k,2}(\mathbb{R}^n)$. We further assume that \mathfrak{a} satisfies the *ellipticity condition*, that is, there exist positive constants $0 < \lambda \leq \Lambda < \infty$ such that

$$\|a_{\alpha,\beta}\|_{L^\infty(\mathbb{R}^n)} \leq \Lambda \quad \text{for all } \alpha, \beta \text{ with } |\alpha| = k = |\beta| \quad (2.8)$$

and

$$\Re \mathfrak{a}(f, f) \geq \lambda \|\nabla^k f\|_{L^2(\mathbb{R}^n)}^2 \quad \text{for all } f \in W^{k,2}(\mathbb{R}^n), \quad (2.9)$$

here and in what follows, $\Re z$ for any $z \in \mathbb{C}$ denotes the *real part* of z . The *2k-order divergence form homogeneous elliptic operator* L_1 with complex bounded measurable coefficients is then defined to be the operator associated with the form \mathfrak{a} .

Let $\omega \in [0, \pi/2]$. Recall that an operator T in the Hilbert space \mathcal{H} is called *m- ω -accretive* if

- (i) the range of the operator $T + I$, $R(T + I)$, is dense in \mathcal{H} ;
- (ii) for all $u \in D(T)$, $|\arg(Tu, u)| \leq \omega$,

where $D(T)$ denotes the *domain* of T and $\arg(Tu, u)$ the *argument* of (Tu, u) . It is known by [33, Proposition 7.1.1] that every closed *m- ω -accretive* operator is of type ω (see [33, p. 173]).

From [6], it follows that L_1 is closed and maximal accretive (see [33, p. 327] for the definition), which further yields that $R(L_1 + I)$ is dense in $L^2(\mathbb{R}^n)$ (see, for example [33, Proposition C.7.2]). Moreover, by the ellipticity condition (2.8) and (2.9), we obtain that for all $f \in W^{k,2}(\mathbb{R}^n)$,

$$|\tan(\arg(L_1 f, f))| = \left| \frac{\Im(L_1 f, f)}{\Re(L_1 f, f)} \right| \leq \frac{\Lambda}{\lambda},$$

here and in what follows, $\Im z$ for any $z \in \mathbb{C}$ denotes the *imaginary part* of z . Thus, $|\arg(L_1 f, f)| \leq \arctan \frac{\Lambda}{\lambda}$, which, together with the fact that $R(L_1 + I)$ is dense in $L^2(\mathbb{R}^n)$, shows that L_1 is an *m- $\arctan \frac{\Lambda}{\lambda}$ -accretive* operator in $L^2(\mathbb{R}^n)$ with the angle $\arctan \frac{\Lambda}{\lambda} \in [\pi/4, \pi/2]$. Thus, L_1 is an operator of type $\arctan \frac{\Lambda}{\lambda}$.

Now, we show that L_1 is one-to-one. Let $N(L_1) := \{f \in D(L_1) : L_1 f = 0\}$ be the *null space* of L_1 . For any fixed $f \in N(L_1)$, by the elliptic condition (2.8) and (2.9), we have

$$\int_{\mathbb{R}^n} |\nabla^k f(x)|^2 dx \sim |(L_1 f, f)| = 0, \quad (2.10)$$

which implies that $\nabla^k f = 0$ almost everywhere in \mathbb{R}^n . In what follows, denote by $C_c^\infty(\mathbb{R}^n)$ the *space of all C^∞ functions with compact support in \mathbb{R}^n* . For all $f \in W^{k,2}(\mathbb{R}^n)$, by the density of $C_c^\infty(\mathbb{R}^n)$ in $W^{k,2}(\mathbb{R}^n)$, there exists a sequence $\{f_j\}_{j \in \mathbb{N}}$ of functions in $C_c^\infty(\mathbb{R}^n)$ such that $\lim_{j \rightarrow \infty} f_j = f$ in $W^{k,2}(\mathbb{R}^n)$. Denote the *Fourier transform* and the *inverse Fourier transform* of f , respectively, by \widehat{f} and f^\vee . If $f \in N(L_1)$, by (2.10), the fact that $f_j \in C_c^\infty(\mathbb{R}^n)$, the multiplication formula of Fourier transform and Plancherel's theorem (see, for example, [28, Theorem 2.2.14]), we have that for all $\varphi \in C_c^\infty(\mathbb{R}^n)$,

$$\begin{aligned} 0 &= (\nabla^k f, \widehat{\varphi}) = \lim_{j \rightarrow \infty} (\nabla^k f_j, \widehat{\varphi}) = \lim_{j \rightarrow \infty} (-1)^k (f_j, \nabla^k \widehat{\varphi}) = \lim_{j \rightarrow \infty} (-1)^k (\widehat{f_j}, (\nabla^k \widehat{\varphi})^\vee) \\ &= \lim_{j \rightarrow \infty} i^k (\widehat{f_j}, (\cdot)^k \varphi(\cdot)) = i^k (\widehat{f}, (\cdot)^k \varphi(\cdot)), \end{aligned}$$

which implies that $\text{supp } \widehat{f} \subset \{0\}$. By [28, Corollary 2.4.2], we have that f is a polynomial, which, together with the fact that $f \in L^2(\mathbb{R}^n)$, implies that $f = 0$. Hence, $N(L_1) = \{0\}$ and L_1 is one-to-one.

Since L_1 is maximal accretive, from [2], it follows that L_1 has a bounded holomorphic functional calculus. Finally, in Proposition 3.4 below, we show that the semigroup $\{e^{-tL_1}\}_{t>0}$ satisfies the k -Davies-Gaffney estimate. Thus, the $2k$ -order divergence form homogenous elliptic operator L_1 with complex bounded measurable coefficients satisfies the assumptions (A_1) , (A_2) and (A_3) .

Let $k \in \mathbb{N}$, $\Delta := \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2}$ be the Laplace operator and $0 \leq V \in L^k_{\text{loc}}(\mathbb{R}^n)$. The $2k$ -order Schrödinger type operator $L_2 := (-\Delta)^k + V^k$ is the associated operator of the following sesquilinear form

$$\mathfrak{b}(f, g) := \int_{\mathbb{R}^n} \nabla^k f(x) \overline{\nabla^k g(x)} dx + \int_{\mathbb{R}^n} [V(x)]^k f(x) \overline{g(x)} dx$$

with domain $D(\mathfrak{b}) := \{f \in W^{k,2}(\mathbb{R}^n) : \int_{\mathbb{R}^n} [V(x)]^k |f(x)|^2 dx < \infty\}$ which is also dense in $L^2(\mathbb{R}^n)$, since $C_c^\infty(\mathbb{R}^n) \subset D(\mathfrak{b})$.

It is easy to see that the $2k$ -order Schrödinger type operator L_2 is a nonnegative self-adjoint operator. From [33], it follows that L_2 is m -0-accretive. Thus, by [33, Proposition 7.1.1], L_2 is a one-to-one operator of type 0. Therefore, L_2 has a bounded H_∞ functional calculus. Moreover, by [9], the semigroup $\{e^{-tL_2}\}_{t>0}$ satisfies a Gaussian type estimate, that is, the integral kernel $e^{-tL_2}(x, y)$ of e^{-tL_2} has the property that there exist positive constants C_2 and C_3 such that for all $t \in (0, \infty)$ and $x, y \in \mathbb{R}^n$,

$$|e^{-tL_2}(x, y)| \leq C_2 t^{-n/(2k)} \exp \left\{ -C_3 \frac{|x-y|^{2k/(2k-1)}}{t^{1/(2k-1)}} \right\},$$

which implies that the semigroup $\{e^{-tL_2}\}_{t>0}$ satisfies the k -Davies-Gaffney estimate immediately. Thus, the $2k$ -order Schrödinger type operator L_2 also satisfies the assumptions (A_1) , (A_2) and (A_3) .

3 k -Davies-Gaffney estimates

In this section, we prove some properties about the k -Davies-Gaffney estimates. We point out that when $k = 1$ and L is a non-negative self-adjoint operator or a second order divergence form elliptic operator with complex bounded measurable coefficients, these properties are already well known (see, for example, [5, 35, 37–39, 41]).

Let $\theta \in [0, \pi/2)$ and E, F be two closed sets in \mathbb{R}^n . A family $\{T(z)\}_{z \in S_\theta^0}$ of operators is said to satisfy the k -Davies-Gaffney estimate in z if there exist positive constants C_4 and C_5 such that for all $f \in L^2(\mathbb{R}^n)$ supported in E and $z \in S_\theta^0$,

$$\|T(z)f\|_{L^2(F)} \leq C_5 \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4 |z|^{1/(2k-1)}} \right\} \|f\|_{L^2(E)}. \quad (3.1)$$

For any operator satisfying the assumptions (A_1) , (A_2) and (A_3) in Section 2, we have the following property.

Lemma 3.1. Assume that the operator L defined in $L^2(\mathbb{R}^n)$ satisfies the assumptions (A_1) , (A_2) and (A_3) in Section 2. Then for all $\ell \in (0, 1)$, $m \in \mathbb{Z}_+$, the family of operators, $\{(zL)^m e^{-zL}\}_{z \in S_{\ell(\frac{\pi}{2}-\omega)}^0}$, satisfies the k -Davies-Gaffney estimate in z , (3.1), with positive constants C_4 and C_5 depending only on $m, \ell, n, k, \omega, \tilde{C}$ and C_1 .

Proof. We prove this lemma by using some ideas from [35]. Since L is of type ω , we know that the semigroup $\{e^{-tL}\}_{t>0}$ can be extended to a holomorphic semigroup $\{e^{-zL}\}_{z \in S_{\pi/2-\omega}^0}$. Thus, for all $z \in S_{\pi/2-\omega}^0$, closed sets $E, F \subset \mathbb{R}^n$ and $f, g \in L^2(\mathbb{R}^n)$ supported respectively in E and F , the function

$$G(z) : z \mapsto (e^{-zL} f, g)$$

is holomorphic on $S_{\pi/2-\omega}^0$. Moreover, G satisfies the following properties:

(i) there exists a nonnegative constant C such that for all $z \in S_{\pi/2-\omega}^0$,

$$|G(z)| \leq C \|f\|_{L^2(E)} \|g\|_{L^2(F)},$$

(ii) there exist nonnegative constants C and C_1 such that for all $t \in (0, \infty)$,

$$|G(t)| \leq C \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_1 t^{1/(2k-1)}} \right\} \|f\|_{L^2(E)} \|g\|_{L^2(F)}.$$

In [47, Lemma 6.18], letting $\psi := \pi/2 - \omega$, $a := C \|f\|_{L^2(E)} \|g\|_{L^2(F)}$, $b := \frac{1}{C_1} [\text{dist}(E, F)]^{2k/(2k-1)}$, $r := t$, $\alpha := \frac{1}{2k-1}$ and $\beta := 0$, we then obtain that for any $z := r e^{i\theta} \in S_{\ell(\pi/2-\omega)}^0$,

$$\begin{aligned} |F(z)| &\lesssim \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{2(2k-1)C_1 r^{1/(2k-1)}} \sin(\pi/2 - \omega - |\theta|) \right\} \|f\|_{L^2(E)} \|g\|_{L^2(F)} \\ &\lesssim \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4 r^{1/(2k-1)}} \right\} \|f\|_{L^2(E)} \|g\|_{L^2(F)}, \end{aligned} \quad (3.2)$$

where $C_4 := \frac{2C_1(2k-1)}{\sin((1-\ell)(\pi/2-\omega))}$. From the analytic property of semigroups and the Cauchy integral formula, it follows that for all $m \in \mathbb{N}$ and $z \in S_{\ell(\pi/2-\omega)}^0$,

$$(zL)^m e^{-zL} = (-z)^m \frac{m!}{2\pi i} \int_{|\xi-z|=\eta|z|} e^{-\xi L} \frac{d\xi}{(\xi-z)^{m+1}}, \quad (3.3)$$

where $\eta \in (0, \sin((1-\ell)(\pi/2-\omega)))$. Thus, for any $z \in S_{\ell(\pi/2-\omega)}^0$, the ball $B(z, \eta|z|) \subset S_{\pi/2-\omega}^0$. Combining (3.2) and (3.3), by Minkowski's inequality, we obtain

$$\begin{aligned} \|(zL)^m e^{-zL} f\|_{L^2(F)} &\lesssim |z|^m \int_{|\xi-z|=\eta|z|} \left| \frac{1}{(\xi-z)^{m+1}} \right| \|e^{-\xi L} f\|_{L^2(F)} |d\xi| \\ &\lesssim \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4 |z|^{1/(2k-1)}} \right\} \|f\|_{L^2(E)}, \end{aligned}$$

which implies that $\{(zL)^m e^{-zL}\}_{S_{\ell(\pi/2-\omega)}^0}$ satisfies the k -Davies-Gaffney estimate in z . This finishes the proof of Lemma 3.1. \square

Lemma 3.2. Let $\{A_t\}_{t>0}$, $\{B_s\}_{s>0}$ be two families of linear operators, C_6 and C_7 two positive constants. Assume that for all closed sets $E, F \subset \mathbb{R}^n$, $f \in L^2(\mathbb{R}^n)$ supported in E and $t \in (0, \infty)$, the following estimates hold:

$$\|A_t f\|_{L^2(F)} \leq C_6 \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_7 t^{1/(2k-1)}} \right\} \|f\|_{L^2(E)}, \quad (3.4)$$

and

$$\|B_s f\|_{L^2(F)} \leq C_6 \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_7 s^{1/(2k-1)}} \right\} \|f\|_{L^2(E)}. \quad (3.5)$$

Then, there exists a positive constant C such that for all $t, s \in (0, \infty)$, all closed sets $E, F \subset \mathbb{R}^n$ and $f \in L^2(\mathbb{R}^n)$ supported in E ,

$$\|A_t B_s f\|_{L^2(F)} \leq C \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{\tilde{C}_7 (\max\{t, s\})^{1/(2k-1)}} \right\} \|f\|_{L^2(E)}, \quad (3.6)$$

where $\tilde{C}_7 := C_7 2^{2k/(2k-1)}$.

Proof. If $\text{dist}(E, F) = 0$, then (3.6) is a simple corollary of (3.4) and (3.5). Now, we assume that $\text{dist}(E, F) > 0$. As in [36], let $\rho := \text{dist}(E, F)$ and $G := \{x \in \mathbb{R}^n : \text{dist}(x, F) < \rho/2\}$. Denote by \overline{G} the closure of G . It is clear that $\text{dist}(E, \overline{G}) \geq \rho/2$. Moreover, by (3.4) and (3.5), we have

$$\begin{aligned} \|A_t(\chi_G B_s f)\|_{L^2(F)} &\leq \|A_t(\chi_G B_s f)\|_{L^2(\mathbb{R}^n)} \lesssim \|B_s f\|_{L^2(\overline{G})} \\ &\lesssim \exp\left\{-\frac{[\text{dist}(E, \overline{G})]^{2k/(2k-1)}}{C_7 s^{1/(2k-1)}}\right\} \|f\|_{L^2(E)} \\ &\sim \exp\left\{-\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_7 2^{2k/(2k-1)} s^{1/(2k-1)}}\right\} \|f\|_{L^2(E)}. \end{aligned}$$

Let $\tilde{C}_7 := C_7 2^{2k/(2k-1)}$. Similarly, by (3.4) and (3.5), we obtain

$$\begin{aligned} \|A_t(\chi_{\mathbb{R}^n \setminus G} B_s f)\|_{L^2(F)} &\lesssim \exp\left\{-\frac{[\text{dist}(\mathbb{R}^n \setminus G, F)]^{2k/(2k-1)}}{C_7 t^{1/(2k-1)}}\right\} \|B_s f\|_{L^2(E)} \\ &\lesssim \exp\left\{-\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{\tilde{C}_7 t^{1/(2k-1)}}\right\} \|f\|_{L^2(E)}. \end{aligned}$$

Combining the above estimates, we have

$$\begin{aligned} \|A_t B_s f\|_{L^2(F)} &\leq \|A_t(\chi_G B_s f)\|_{L^2(F)} + \|A_t(\chi_{\mathbb{R}^n \setminus G} B_s f)\|_{L^2(F)} \\ &\lesssim \left[\exp\left\{-\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{\tilde{C}_7 s^{1/(2k-1)}}\right\} + \exp\left\{-\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{\tilde{C}_7 t^{1/(2k-1)}}\right\} \right] \|f\|_{L^2(E)} \\ &\lesssim \exp\left\{-\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{\tilde{C}_7 \max\{t, s\}^{1/(2k-1)}}\right\} \|f\|_{L^2(E)}, \end{aligned}$$

which completes the proof of Lemma 3.2. \square

Let $\sigma \in [0, \infty)$. As in [39], a family $\{T_t\}_{t>0}$ of operators is said to satisfy the k -Davies-Gaffney estimate of order σ , if there exists a positive constant C_8 , depending on σ , such that for all closed sets $E, F \subset \mathbb{R}^n$, $g \in L^2(\mathbb{R}^n)$ supported in E and $t \in (0, \infty)$,

$$\|T_t g\|_{L^2(F)} \leq C_8 \min\left\{1, \frac{t}{[\text{dist}(E, F)]^{2k}}\right\}^\sigma \|g\|_{L^2(E)}. \quad (3.7)$$

Lemma 3.3. Let $\mu \in (\omega, \pi/2)$, $\psi \in \Psi_{\sigma, \tau}(S_\mu^0)$ for some $\sigma \in (0, \infty)$, $\tau \in (1, \infty)$, and $f \in H_\infty(S_\mu^0)$. Then the family $\{\psi(tL)f(L)\}_{t>0}$ of operators satisfies the k -Davies-Gaffney estimate of order σ , (3.7), with the positive constant C_8 controlled by $\|f\|_{L^\infty(S_\mu^0)}$.

Proof. For any fixed $\psi \in \Psi_{\sigma, \tau}(S_\mu^0) \subset \Psi(S_\mu^0)$ and $f \in H_\infty(S_\mu^0)$, by (2.4) and (2.5), we have

$$\psi(tL)f(L) = \int_{\Gamma_+} e^{-zL} \eta_+(z) dz + \int_{\Gamma_-} e^{-zL} \eta_-(z) dz, \quad (3.8)$$

where $\Gamma_\pm := \mathbb{R}^+ e^{\pm i(\pi/2 - \theta)}$ and for all $z \in \Gamma_\pm$,

$$\eta_\pm(z) = \frac{1}{2\pi i} \int_{\gamma_\pm} e^{\xi z} \psi(t\xi) f(\xi) d\xi,$$

$\gamma_\pm := \mathbb{R}^+ e^{\pm i\nu}$ and $0 \leq \omega < \theta < \nu < \mu < \pi/2$. It was proved in [39, (2.32)] that for all $z \in \Gamma_\pm$,

$$|\eta_\pm(z)| \lesssim \frac{\|f\|_{L^\infty(S_\mu^0)}}{t} \min\left\{1, \left(\frac{t}{|z|}\right)^{\sigma+1}\right\}. \quad (3.9)$$

Thus, by (3.8) and Minkowski's inequality, we have that for all $g \in L^2(\mathbb{R}^n)$ supported in E ,

$$\|\psi(tL)f(L)g\|_{L^2(F)} \leq \int_{\Gamma_+} \|e^{-zL}g\|_{L^2(F)} |\eta_+(z)| |dz| + \int_{\Gamma_-} \|e^{-zL}g\|_{L^2(F)} |\eta_-(z)| |dz| =: J_+ + J_-.$$

Since $\pi/2 - \theta < \pi/2 - \omega$, there exists a positive number $\ell \in (0, 1)$ such that $\pi/2 - \theta < \ell(\pi/2 - \omega)$, which immediately yields that $S_{\pi/2-\theta}^0 \subset S_{\ell(\pi/2-\omega)}^0$. Thus, by Lemma 3.1, the family $\{e^{-zL}\}_{z \in S_{\pi/2-\theta}^0}$ satisfies the k -Davies-Gaffney estimate in z , which, together with (3.9), implies that

$$\begin{aligned} J_{\pm} &\lesssim \|g\|_{L^2(E)} \int_{\Gamma_{\pm}} \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4|z|^{1/(2k-1)}} \right\} |\eta_{\pm}(z)| |dz| \\ &\lesssim \|f\|_{L^\infty(S_\mu^0)} \|g\|_{L^2(E)} \int_{\Gamma_{\pm}} \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4|z|^{1/(2k-1)}} \right\} \min \left\{ 1, \left(\frac{t}{|z|} \right)^{\sigma+1} \right\} \frac{1}{t} |dz| \\ &\lesssim \left[\int_{\{z \in \Gamma_{\pm}: |z| \leq t\}} \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4|z|^{1/(2k-1)}} \right\} \min \left\{ 1, \left(\frac{t}{|z|} \right)^{\sigma+1} \right\} \frac{1}{t} |dz| \right. \\ &\quad \left. + \int_{\{z \in \Gamma_{\pm}: |z| > t\}} \cdots \right] \|f\|_{L^\infty(S_\mu^0)} \|g\|_{L^2(E)} \\ &=: [O_1 + O_2] \|f\|_{L^\infty(S_\mu^0)} \|g\|_{L^2(E)}. \end{aligned}$$

We estimate O_1 by

$$O_1 \lesssim \int_{\{z \in \Gamma_{\pm}: |z| \leq t\}} \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4|z|^{1/(2k-1)}} \right\} \frac{1}{t} |dz| \lesssim \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4|t|^{1/(2k-1)}} \right\}.$$

On the other hand, O_2 can be written into

$$O_2 \lesssim \int_{\{z \in \Gamma_{\pm}: |z| > t\}} \exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_4|z|^{1/(2k-1)}} \right\} \left(\frac{t}{|z|} \right)^{\sigma+1} \frac{1}{t} |dz|.$$

If $t \geq [\text{dist}(E, F)]^{2k}$, in this case, we have

$$O_2 \lesssim \int_{\{z \in \Gamma_{\pm}: |z| > t\}} \left(\frac{t}{|z|} \right)^{\sigma+1} \frac{1}{t} |dz| \sim 1.$$

If $t < [\text{dist}(E, F)]^{2k}$, by choosing $N \in [\sigma, \infty)$, we obtain

$$\begin{aligned} O_2 &\lesssim \int_{\{z \in \Gamma_{\pm}: t < |z| \leq [\text{dist}(E, F)]^{2k}\}} \left(\frac{|z|}{[\text{dist}(E, F)]^{2k}} \right)^N \left(\frac{t}{|z|} \right)^{\sigma+1} \frac{1}{t} |dz| \\ &\quad + \int_{\{z \in \Gamma_{\pm}: |z| > [\text{dist}(E, F)]^{2k}\}} \left(\frac{t}{|z|} \right)^{\sigma+1} \frac{1}{t} |dz| \\ &\lesssim \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^N + \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^{\sigma} \sim \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^{\sigma}. \end{aligned}$$

Combining the estimates of O_1 and O_2 , we obtain that $\{\psi(tL)f(L)\}_{t>0}$ satisfies the k -Davies-Gaffney estimate of order σ , which completes the proof of Lemma 3.3. \square

Now, we turn to some properties of the operators L_1 and L_2 given in Section 2. First, we introduce the definition of the Legendre transform. Let h be a real valued function defined on $[0, \infty)$. The *Legendre transform* h^\sharp of h is defined by setting, for all $s \in \mathbb{R}$,

$$h^\sharp(s) := \sup_{t \geq 0} \{st - h(t)\}. \quad (3.10)$$

We have the following proposition about the operator L_1 .

Proposition 3.4. *Let L_1 be the $2k$ -order divergence form homogeneous elliptic operator defined as in Section 2. Then, the semigroup $\{e^{-tL_1}\}_{t>0}$ satisfies the k -Davies-Gaffney estimate.*

Proof. We prove Proposition 3.4 by borrowing some ideas from [10, 11, 19]. In [11, Theorem 1.2], letting $(\Omega, \mathcal{U}, \mu, d)$ be the usual Euclidean space \mathbb{R}^n , endowed with the Lebesgue measure dx and the Euclidean distant d , and with the set class \mathcal{U} being the set of all Lebesgue measurable sets, and also letting

$$\mathcal{A} := \{\phi \in C^\infty(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n) : \|D^\alpha \phi\|_{L^\infty(\mathbb{R}^n)} \leq 1, 1 \leq |\alpha| \leq k\},$$

$p \equiv q := 2$, $\alpha \equiv \beta \equiv \gamma := 0$, $r := t^{1/(2k)}$, $h(x) := x^{2k/(2k-1)}$ for all $x \in [0, \infty)$ and $\mathcal{R} := e^{-tL_1}$, we then obtain the following two equivalent statements:

- (i) There exists a positive constant $C(k)$, depending on k , such that for all $\phi \in \mathcal{A}$, $\rho \in [0, \infty)$ and $t \in (0, \infty)$,

$$\|e^{-\rho\phi} e^{-tL_1} e^{\rho\phi}\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq e^{C(k)h^\sharp(\rho t^{1/(2k)})}, \quad (3.11)$$

where, by (3.10),

$$h^\sharp(\rho t^{1/(2k)}) := \sup_{s \geq 0} \left\{ \rho t^{1/(2k)} s - s^{2k/(2k-1)} \right\} = \left[\frac{(2k-1)^{2k-1}}{(2k)^{2k}} \right] \rho^{2k} t;$$

- (ii) There exists a positive constant C_1 such that for all the closed sets E and F of \mathbb{R}^n and $t \in (0, \infty)$,

$$\|\chi_E e^{-tL_1} \chi_F\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq \exp \left(-\frac{\text{dist}(E, F)}{C_1 t^{1/(2k)}} \right)^{2k/(2k-1)}. \quad (3.12)$$

Recall that in this case, by [19, Lemma 4], $d(E, F)$ defined in [11, (1.4)] is equivalent to $\inf_{x \in E, y \in F} |x - y|$ and, moreover, $C(k)$ in [11, Theorem 1.2(ii)] is assumed to be 1. However, by the change of variables, we can easily see that the above equivalent statements are a directly corollary of the equivalence of (ii) and (iii) of Theorem 1.2 in [11].

Notice that by the density of the simple functions in $L^2(\mathbb{R}^n)$, (3.12) is equivalent to the k -Davies-Gaffney estimate. Thus, to prove that the semigroup $\{e^{-tL_1}\}_{t>0}$ satisfies the k -Davies-Gaffney estimate, by the above equivalence of (i) and (ii), it suffices to prove (3.11).

To this end, let \mathfrak{a} be the sesquilinear form as in (2.7) associated with L_1 . Recall that its *twisted form* is defined by setting, for all $\rho \in [0, \infty)$, $\phi \in \mathcal{A}$ and $f, g \in W^{k,2}(\mathbb{R}^n)$,

$$\mathfrak{a}_{\rho\phi}(f, g) := \mathfrak{a}(e^{\rho\phi} f, e^{-\rho\phi} g),$$

which, together with the Leibniz formula, further yields that there exist positive constants $C(\alpha, \gamma)$ and $C(\beta, \tilde{\gamma})$ with $|\alpha| = |\beta| = k$, $0 < \gamma \leq \alpha$ and $0 < \tilde{\gamma} \leq \beta$,

$$\begin{aligned} \mathfrak{a}_{\rho\phi}(f, f) &= \sum_{|\alpha|=|\beta|=k} \int_{\mathbb{R}^n} a_{\alpha,\beta}(x) \partial^\alpha (e^{\rho\phi} f)(x) \overline{\partial^\beta (e^{-\rho\phi} f)(x)} dx \\ &= \sum_{|\alpha|=|\beta|=k} \int_{\mathbb{R}^n} a_{\alpha,\beta}(x) \left\{ \left[\sum_{0 < \gamma \leq \alpha} C(\alpha, \gamma) \partial^\gamma e^{\rho\phi(x)} \partial^{\alpha-\gamma} f(x) + e^{\rho\phi(x)} \partial^\alpha f(x) \right] \right. \\ &\quad \left. + \left[\sum_{0 < \tilde{\gamma} \leq \beta} C(\beta, \tilde{\gamma}) \overline{\partial^{\tilde{\gamma}} e^{-\rho\phi(x)} \partial^{\beta-\tilde{\gamma}} f(x)} + \overline{e^{-\rho\phi(x)} \partial^\beta f(x)} \right] \right\} dx \\ &= \sum_{|\alpha|=|\beta|=k} \int_{\mathbb{R}^n} a_{\alpha,\beta}(x) \left\{ \left[\sum_{0 < \gamma \leq \alpha, 0 < \tilde{\gamma} \leq \beta} C(\alpha, \gamma) C(\beta, \tilde{\gamma}) \partial^\gamma e^{\rho\phi(x)} \partial^{\alpha-\gamma} f(x) \overline{\partial^{\tilde{\gamma}} e^{-\rho\phi(x)} \partial^{\beta-\tilde{\gamma}} f(x)} \right] \right\} \end{aligned}$$

$$+ \left[\frac{\overline{e^{-\rho\phi(x)}\partial^\beta f(x)}}{\sum_{0 < \gamma \leq \alpha} C(\alpha, \gamma) \partial^\gamma e^{\rho\phi(x)} \partial^{\alpha-\gamma} f(x)} \right] \\ + \left[\frac{\overline{e^{\rho\phi(x)}\partial^\alpha f(x)}}{\sum_{0 < \tilde{\gamma} \leq \beta} C(\beta, \tilde{\gamma}) \partial^{\tilde{\gamma}} e^{-\rho\phi(x)} \partial^{\beta-\tilde{\gamma}} f(x)} \right] \Bigg\} dx + \mathfrak{a}(f, f).$$

Let $C(\alpha, 0) := 1 =: C(\beta, 0)$ and $\tilde{C}(k) := \Lambda \sum_{|\alpha|=|\beta|=k} [\sum_{0 \leq \gamma \leq \alpha, 0 \leq \tilde{\gamma} \leq \beta} C(\alpha, \gamma) C(\beta, \tilde{\gamma})]$, where Λ is as in (2.8). By this estimate, $\phi \in \mathcal{A}$, (2.8) and Hölder's inequality, we further have

$$|\mathfrak{a}_{\rho, \phi}(f, f) - \mathfrak{a}(f, f)| \\ \leq \Lambda \sum_{|\alpha|=|\beta|=k} \int_{\mathbb{R}^n} \left\{ \sum_{0 \leq \gamma \leq \alpha, 0 \leq \tilde{\gamma} \leq \beta} C(\alpha, \gamma) C(\beta, \tilde{\gamma}) \left| \rho^{|\gamma|} \partial^{\alpha-\gamma} f(x) \rho^{|\tilde{\gamma}|} \overline{\partial^{\beta-\tilde{\gamma}} f(x)} \right| \right\} dx \\ \leq \Lambda \sum_{|\alpha|=|\beta|=k} \sum_{0 \leq \gamma \leq \alpha, 0 \leq \tilde{\gamma} \leq \beta} C(\alpha, \gamma) C(\beta, \tilde{\gamma}) \left\{ \int_{\mathbb{R}^n} \left| \rho^{|\gamma|} \partial^{\alpha-\gamma} f(x) \right|^2 dx \right\}^{1/2} \left\{ \int_{\mathbb{R}^n} \left| \rho^{|\tilde{\gamma}|} \partial^{\beta-\tilde{\gamma}} f(x) \right|^2 dx \right\}^{1/2} \\ =: \Lambda \sum_{|\alpha|=|\beta|=k} \sum_{0 \leq \gamma \leq \alpha, 0 \leq \tilde{\gamma} \leq \beta} C(\alpha, \gamma) C(\beta, \tilde{\gamma}) \mathbf{I}_1 \times \mathbf{I}_2.$$

Applying Plancherel's theorem, (2.9) and Young's inequality with $\epsilon \in (0, \frac{\lambda}{4C(k)})$, we obtain that there exists a positive constant $C(\epsilon)$ such that for all $\tilde{\lambda} \in (C(\epsilon)\tilde{C}(k), \infty)$,

$$(\mathbf{I}_1)^2 \leq \int_{\mathbb{R}^n} \left[\rho^{|\gamma|} |\xi|^{k-|\gamma|} \left| \widehat{f}(\xi) \right| \right]^2 d\xi \leq \int_{\mathbb{R}^n} [C(\epsilon)\rho^{2k} + \epsilon|\xi|^{2k}] \left| \widehat{f}(\xi) \right|^2 d\xi \\ \leq C(\epsilon)\rho^{2k} \|f\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|\nabla^k f\|_{L^2(\mathbb{R}^n)}^2 \leq C(\epsilon)\rho^{2k} \|f\|_{L^2(\mathbb{R}^n)}^2 + \frac{\epsilon}{\tilde{\lambda}} \Re \mathfrak{a}(f, f),$$

which, together with a similar estimate for \mathbf{I}_2 , shows that

$$|\mathfrak{a}_{\rho\phi}(f, f) - \mathfrak{a}(f, f)| \leq \frac{1}{4} \Re \mathfrak{a}(f, f) + \tilde{\lambda} \rho^{2k} \|f\|_{L^2(\mathbb{R}^n)}^2. \quad (3.13)$$

Denote by $L_{\rho\phi}(=e^{-\rho\phi}L_1e^{\rho\phi})$ the operator associated with $\mathfrak{a}_{\rho\phi}$. Let $f_t := e^{-tL_{\rho\phi}}f$. By (3.13), we have

$$\frac{d}{dt} \|f_t\|_{L^2(\mathbb{R}^n)}^2 = -(L_{\rho\phi}f_t, f_t) - (f_t, L_{\rho\phi}f_t) = -2\Re \mathfrak{a}_{\rho\phi}(f_t, f_t) \\ = 2[\Re(\mathfrak{a}(f_t, f_t) - \mathfrak{a}_{\rho\phi}(f_t, f_t)) - \Re \mathfrak{a}(f_t, f_t)] \\ \leq 2|\mathfrak{a}_{\rho\phi}(f_t, f_t) - \mathfrak{a}(f_t, f_t)| - 2\Re \mathfrak{a}(f_t, f_t) \\ \leq \frac{1}{2} \Re \mathfrak{a}(f_t, f_t) + 2\tilde{\lambda} \rho^{2k} \|f_t\|_{L^2(\mathbb{R}^n)}^2 - 2\Re \mathfrak{a}(f_t, f_t) \leq 2\tilde{\lambda} \rho^{2k} \|f_t\|_{L^2(\mathbb{R}^n)}^2.$$

Thus,

$$\|f_t\|_{L^2(\mathbb{R}^n)}^2 = \|e^{-tL_{\rho\phi}}f\|_{L^2(\mathbb{R}^n)}^2 \leq \exp\{2\tilde{\lambda}\rho^{2k}t\} \|f\|_{L^2(\mathbb{R}^n)}^2 \leq \exp\left\{2C(k)h^\sharp(\rho t^{1/(2k)})\right\} \|f\|_{L^2(\mathbb{R}^n)}^2.$$

That is, (3.11) holds. Therefore, $\{e^{-tL_1}\}_{t>0}$ satisfies the k -Davies-Gaffney estimate, which completes the proof of Proposition 3.4. \square

Remark 3.5. In the proof of Proposition 3.4, we obtain the estimate (3.11) by following the proof of [19, Lemma 2]. The same method should also work for the proof of [10, Proposition 3.1]. Notice that the scaling method mentioned in the last two lines of [10, p. 143] may not be valid when used to remove the factor $e^{(\alpha w + \epsilon)t}$ appearing in the proof of [10, Proposition 3.1], as the authors claimed therein.

We also have the following gradient estimate for L_1 and L_2 .

Proposition 3.6. Let $k \in \mathbb{N}$, L_1 be the $2k$ -order divergence form elliptic operator and L_2 the $2k$ -order Schrödinger type operator defined as in Section 2. Then, $\{\sqrt{t}\nabla^k e^{-tL_i}\}_{t>0}$ for $i \in \{1, 2\}$ satisfies the k -Davies-Gaffney estimate.

Proof. For any Hilbert space \mathcal{H} , let $(\cdot, \cdot)_{\mathcal{H}}$ be the inner product of \mathcal{H} . By Hölder's inequality and the fact that $\{tL_i e^{-tL_i}\}_{t>0}$ and $\{e^{-tL_i}\}_{t>0}$ satisfy the k -Davies-Gaffney estimate which are deduced respectively from Proposition 3.4 and Lemma 3.1, we conclude that for all closed sets $E, F \subset \mathbb{R}^n$, $f \in L^2(\mathbb{R}^n)$ supported in E , and $t \in (0, \infty)$,

$$\begin{aligned} \left\| \sqrt{t}\nabla^k e^{-tL_i} f \right\|_{L^2(F)}^2 &\lesssim \left| (tL_i e^{-tL_i} f, e^{-tL_i} f)_{L^2(F)} \right| \\ &\lesssim \|tL_i e^{-tL_i} f\|_{L^2(F)} \|e^{-tL_i} f\|_{L^2(F)} \\ &\lesssim \left(\exp \left\{ -\frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{C_1 t^{1/(2k-1)}} \right\} \right)^2 \|f\|_{L^2(E)}^2, \end{aligned}$$

which implies that $\{\sqrt{t}\nabla^k e^{-tL_i}\}_{t>0}$ also satisfies the k -Davies-Gaffney estimate. This finishes the proof of Proposition 3.6. \square

4 Molecular characterizations of $H_L^p(\mathbb{R}^n)$

Assume that the operator L satisfies the assumptions (A_1) , (A_2) and (A_3) in Section 2. In this section, we introduce the Hardy space $H_L^p(\mathbb{R}^n)$ by means of the L -adapted square function and characterize these Hardy spaces by the molecular decomposition. First, we recall some notions.

Let

$$\Gamma(x) := \{(y, t) \in \mathbb{R}^n \times (0, \infty) : |x - y| < t\}$$

be the cone with vertex $x \in \mathbb{R}^n$. For all $f \in L^2(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$, the L -adapted square function $S_L f$ is defined by

$$S_L f(x) := \left\{ \iint_{\Gamma(x)} |t^{2k} L e^{-t^{2k} L} f(y)|^2 \frac{dy dt}{t^{n+1}} \right\}^{1/2}. \quad (4.1)$$

Definition 4.1. Let $p \in (0, 1]$ and L satisfy the assumptions (A_1) , (A_2) and (A_3) in Section 2. A function $f \in L^2(\mathbb{R}^n)$ is said to be in $\mathbb{H}_L^p(\mathbb{R}^n)$ if $S_L f \in L^p(\mathbb{R}^n)$; moreover, define $\|f\|_{\mathbb{H}_L^p(\mathbb{R}^n)} := \|S_L f\|_{L^p(\mathbb{R}^n)}$. The Hardy space $H_L^p(\mathbb{R}^n)$ is then defined to be the completion of $\mathbb{H}_L^p(\mathbb{R}^n)$ with respect to the quasi-norm $\|\cdot\|_{H_L^p(\mathbb{R}^n)}$.

Remark 4.2. Since both the $2k$ -order divergence form homogenous elliptic operator L_1 with complex bounded measurable coefficients and the $2k$ -order Schrödinger type operator L_2 satisfy the assumptions (A_1) , (A_2) and (A_3) , we then define the Hardy spaces $H_{L_1}^p(\mathbb{R}^n)$ and $H_{L_2}^p(\mathbb{R}^n)$, respectively, associated with L_1 and L_2 as in Definition 4.1. In particular, when $k = 1$, $H_{L_1}^p(\mathbb{R}^n)$ is just the Hardy space $H_{-\text{div}(A\nabla)}^p(\mathbb{R}^n)$ associated with the second order divergence form elliptic operator $-\text{div}(A\nabla)$ with complex bounded measurable coefficients in [37–39, 43] and $H_{L_2}^1(\mathbb{R}^n)$ appears in [24, 25, 35, 56]; when $k = 2$, $H_{L_2}^1(\mathbb{R}^n)$ was also studied in [12].

In what follows, a *cube* always means a closed cube whose sides are parallel to the coordinate axes. Let $Q \subset \mathbb{R}^n$ be a cube with the side length $l(Q)$. For $i \in \mathbb{Z}_+$, denote by $S_i(Q)$ the *dyadic annuli* based on Q , namely, $S_0(Q) := Q$ and $S_i(Q) := 2^i Q \setminus (2^{i-1} Q)$ for $i \in \mathbb{N}$, where $2^i Q$ is the cube with the same center as Q and the side length $2^i l(Q)$.

Definition 4.3. Let $p \in (0, 1]$, $\epsilon \in (0, \infty)$, $M \in \mathbb{N}$ and L satisfy the assumptions (A_1) , (A_2) and (A_3) in Section 2. A function $m \in L^2(\mathbb{R}^n)$ is called an (H_L^p, ϵ, M) -molecule if there exists a cube $Q \subset \mathbb{R}^n$ such that

- (i) for each $\ell \in \{1, \dots, M\}$, m belongs to the range of L^ℓ in $L^2(\mathbb{R}^n)$;
(ii) for all $i \in \mathbb{Z}_+$ and $\ell \in \{0, 1, \dots, M\}$,

$$\left\| ([l(Q)]^{-2k} L^{-1})^\ell m \right\|_{L^2(S_i(Q))} \leq [2^i l(Q)]^{n(\frac{1}{2} - \frac{1}{p})} 2^{-i\epsilon}. \quad (4.2)$$

Assume that $\{m_j\}_{j=0}^\infty$ is a sequence of (H_L^p, ϵ, M) -molecules and $\{\lambda_j\}_{j=0}^\infty \in l^p$. For any $f \in L^2(\mathbb{R}^n)$, if $f = \sum_{j=0}^\infty \lambda_j m_j$ in $L^2(\mathbb{R}^n)$, then $\sum_{j=0}^\infty \lambda_j m_j$ is called a *molecular* $(H_L^p, 2, \epsilon, M)$ -representation of f .

We now introduce the notion of a molecular Hardy space $H_{L, \text{mol}, M}^p(\mathbb{R}^n)$ generated by (H_L^p, ϵ, M) -molecules.

Definition 4.4. Let $p \in (0, 1]$, $\epsilon \in (0, \infty)$, $M \in \mathbb{N}$ and L satisfy the assumptions (A_1) , (A_2) and (A_3) in Section 2. The *molecular Hardy space* $H_{L, \text{mol}, M}^p(\mathbb{R}^n)$ is defined to be the completion of the space

$$\mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n) := \{f : f \text{ has a molecular } (H_L^p, 2, \epsilon, M) \text{ - representation}\}$$

with respect to the *quasi-norm*

$$\|f\|_{H_{L, \text{mol}, M}^p(\mathbb{R}^n)} := \inf \left\{ \left(\sum_{j=0}^\infty |\lambda_j|^p \right)^{1/p} : f = \sum_{j=0}^\infty \lambda_j m_j \text{ is a molecular } (H_L^p, 2, \epsilon, M) \text{ - representation} \right\},$$

where the infimum is taken over all the molecular $(H_L^p, 2, \epsilon, M)$ -representations of f as above.

Now, we establish the molecular characterization of the Hardy space $H_L^p(\mathbb{R}^n)$.

Theorem 4.5. Let $p \in (0, 1]$, $\epsilon \in (0, \infty)$, L satisfy the assumptions (A_1) , (A_2) and (A_3) in Section 2 and $M \in \mathbb{N}$ such that $M > \frac{n}{2k}(\frac{1}{p} - \frac{1}{2})$. Then, $H_L^p(\mathbb{R}^n) = H_{L, \text{mol}, M}^p(\mathbb{R}^n)$ with equivalent norms.

To prove Theorem 4.5, by Definitions 4.1 and 4.4, it suffices to prove that

$$\mathbb{H}_L^p(\mathbb{R}^n) = \mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n), \quad M > \frac{n}{2k} \left(\frac{1}{p} - \frac{1}{2} \right),$$

with equivalent norms. We divide the proof into two parts: (i) $\mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n) \subset \mathbb{H}_L^p(\mathbb{R}^n)$; (ii) $\mathbb{H}_L^p(\mathbb{R}^n) \subset \mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n)$.

To prove the inclusion $\mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n) \subset \mathbb{H}_L^p(\mathbb{R}^n)$, we need the following key lemma which is just [39, Lemma 3.8]. Recall that a *nonnegative sublinear operator* T means that T is sublinear and $Tf \geq 0$ for all f in the domain of T .

Lemma 4.6. Let $p \in (0, 1]$, $M \in \mathbb{N}$ and T be a linear operator, or a nonnegative sublinear operator, which is of weak type $(2, 2)$, that is, there exists a positive constant C such that for all $\eta \in (0, \infty)$ and $f \in L^2(\mathbb{R}^n)$,

$$|\{x \in \mathbb{R}^n : |Tf(x)| > \eta\}| \leq C\eta^{-2} \|f\|_{L^2(\mathbb{R}^n)}^2.$$

Assume that there exists a positive constant C such that for all (H_L^p, ϵ, M) -molecules m , $\|Tm\|_{L^p(\mathbb{R}^n)} \leq C$. Then the operator T is bounded from $H_{L, \text{mol}, M}^p(\mathbb{R}^n)$ to $L^p(\mathbb{R}^n)$.

Proof of Theorem 4.5: the inclusion $\mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n) \subset \mathbb{H}_L^p(\mathbb{R}^n)$. Recall that L is a one-to-one operator of type ω having a bounded H_∞ functional calculus. For all $x \in \mathbb{R}^n$, $\psi \in \Psi(S_\mu^0)$ defined as in Section 2, set $\psi_t(x) := \psi(tx)$ for all $t \in (0, \infty)$. The *quadratic norm* $\|g\|_{T, \psi}$, associated with the operator L in $L^2(\mathbb{R}^n)$ and ψ , is defined by

$$\|g\|_{T, \psi} := \left\{ \int_0^\infty \|\psi_t(T)g\|_{L^2(\mathbb{R}^n)}^2 \frac{dt}{t} \right\}^{1/2}$$

for all $g \in [L^2(\mathbb{R}^n)]_{T,\psi}$ which is a subspace of $L^2(\mathbb{R}^n)$ such that the above integral is finite. Since L has a bounded H_∞ functional calculus on $L^2(\mathbb{R}^n)$, it follows from [2] that $[L^2(\mathbb{R}^n)]_{T,\psi} = L^2(\mathbb{R}^n)$ and for all $g \in L^2(\mathbb{R}^n)$,

$$\|g\|_{T,\psi} \lesssim \|g\|_{L^2(\mathbb{R}^n)}. \quad (4.3)$$

By Fubini's theorem, we have that $\|S_L g\|_{L^2(\mathbb{R}^n)} \sim \|g\|_{T,\psi_0}$, where $\psi_0(z) := ze^{-z} \in \Psi(S_\mu^0)$ for all $\mu \in (0, \pi/2)$. Thus, S_L is bounded on $L^2(\mathbb{R}^n)$. By Lemma 4.6, to prove the inclusion $\mathbb{H}_{L,\text{mol},M}^p(\mathbb{R}^n) \subset \mathbb{H}_L^p(\mathbb{R}^n)$, it suffices to prove that for all (H_L^p, ϵ, M) -molecules m with $M > \frac{n}{2k}(\frac{1}{p} - \frac{1}{2})$,

$$\|S_L m\|_{L^p(\mathbb{R}^n)} \lesssim 1. \quad (4.4)$$

Let Q be the cube associated with m as in Definition 4.3. Let $j_0 \in \mathbb{N}$ be such that $2^{j_0-1} < \sqrt{n} \leq 2^{j_0}$. By Minkowski's inequality, Hölder's inequality and the $L^2(\mathbb{R}^n)$ -boundedness of S_L , we see that

$$\begin{aligned} \|S_L m\|_{L^p(\mathbb{R}^n)} &\leq \|S_L m\|_{L^p(2^{j_0+4}Q)} + \sum_{j=j_0+5}^{\infty} \|S_L m\|_{L^p(S_j(Q))} \\ &\lesssim \|m\|_{L^2(\mathbb{R}^n)} |2^{j_0+4}Q|^{\frac{1}{p}-\frac{1}{2}} + \sum_{j=j_0+5}^{\infty} \|S_L m\|_{L^2(S_j(Q))} |S_j(Q)|^{\frac{1}{p}-\frac{1}{2}}. \end{aligned}$$

For $\|m\|_{L^2(\mathbb{R}^n)}$, from Minkowski's inequality and the size condition (4.2) of m , it follows that

$$\|m\|_{L^2(\mathbb{R}^n)} \leq \sum_{j=0}^{\infty} \|m\|_{L^2(S_j(Q))} \leq \sum_{j=0}^{\infty} [2^j l(Q)]^{n(\frac{1}{2}-\frac{1}{p})} 2^{-j\epsilon} \lesssim [l(Q)]^{n(\frac{1}{2}-\frac{1}{p})}. \quad (4.5)$$

For $j \in \{j_0+5, \dots\}$, let $I_j := \|S_L m\|_{L^2(S_j(Q))}$. Then,

$$\begin{aligned} (I_j)^2 &= \int_{S_j(Q)} |S_L m|^2 dx = \int_{S_j(Q)} \int_0^\infty \int_{|y-x|<t} |t^{2k} L e^{-t^{2k}L} m(y)|^2 \frac{dy dt}{t^{n+1}} dx \\ &= \int_{S_j(Q)} \int_0^{2^{\theta(j-5)}l(Q)} \int_{|y-x|<t} |t^{2k} L e^{-t^{2k}L} m(y)|^2 \frac{dy dt}{t^{n+1}} dx + \int_{S_j(Q)} \int_{2^{\theta(j-5)}l(Q)}^\infty \int_{|y-x|<t} \dots \\ &=: B_j + D_j, \end{aligned}$$

where $\theta \in (0, 1)$ is determined later.

For D_j , let $b := L^{-M}m$. By Fubini's theorem, Lemma 3.1 and the size condition (4.2) of m , we conclude that

$$\begin{aligned} D_j &= \int_{S_j(Q)} \int_{2^{\theta(j-5)}l(Q)}^\infty \int_{|y-x|<t} |t^{2k} L e^{-t^{2k}L} L^M b(y)|^2 \frac{dy dt}{t^{n+1}} dx \\ &= \int_{S_j(Q)} \int_{2^{\theta(j-5)}l(Q)}^\infty \int_{|y-x|<t} |t^{2k(M+1)} L^{M+1} e^{-t^{2k}L} b(y)|^2 \frac{dy dt}{t^{n+1+4kM}} dx \\ &\lesssim \int_{2^{\theta(j-5)}l(Q)}^\infty \left\| t^{2k(M+1)} L^{M+1} e^{-t^{2k}L} b \right\|_{L^2(\mathbb{R}^n)}^2 \frac{dt}{t^{4kM+1}} \lesssim \int_{2^{\theta(j-5)}l(Q)}^\infty \|b\|_{L^2(\mathbb{R}^n)}^2 \frac{dt}{t^{4kM+1}} \\ &\sim \|b\|_{L^2(\mathbb{R}^n)}^2 \left[2^{\theta(j-5)}l(Q) \right]^{-4kM} \sim \left[\sum_{i=0}^\infty \|b\|_{L^2(S_i(Q))}^2 \right] \left[2^{\theta(j-5)}l(Q) \right]^{-4kM} \\ &\lesssim [l(Q)]^{4kM+2n(\frac{1}{2}-\frac{1}{p})} \left[2^{\theta(j-5)}l(Q) \right]^{-4kM} \sim 2^{-j[4kM\theta+n(1-\frac{2}{p})]} [2^j l(Q)]^{n(1-\frac{2}{p})}. \end{aligned}$$

Recall that $M > \frac{n}{2k}(1/p - 1/2)$. Letting θ be sufficiently close to 1 such that $\alpha_0 := 2kM\theta - n(1/p - 1/2) > 0$, we then obtain

$$D_j \lesssim 2^{-2j\alpha_0} |S_j(Q)|^{1-\frac{2}{p}}. \quad (4.6)$$

To estimate B_j , let $\tilde{S}_j(Q) := 2^{j+j_0+1}Q \setminus (2^{j-j_0-2}Q)$ and $\hat{S}_j(Q) := 2^{j+j_0+2}Q \setminus (2^{j-j_0-3}Q)$. By Fubini's theorem, we see that

$$\begin{aligned} B_j &\lesssim \int_0^{2^{\theta(j-5)l(Q)}} \int_{\tilde{S}_j(Q)} \left| t^{2k} L e^{-t^{2k}L} m(y) \right|^2 \frac{dy dt}{t} \\ &\lesssim \int_0^{2^{\theta(j-5)l(Q)}} \int_{\tilde{S}_j(Q)} \left| t^{2k} L e^{-t^{2k}L} (\chi_{2^{j-j_0-3}Q} m)(y) \right|^2 \frac{dy dt}{t} \\ &\quad + \int_0^{2^{\theta(j-5)l(Q)}} \int_{\tilde{S}_j(Q)} \left| t^{2k} L e^{-t^{2k}L} (\chi_{\hat{S}_j(Q)} m)(y) \right|^2 \frac{dy dt}{t} \\ &\quad + \int_0^{2^{\theta(j-5)l(Q)}} \int_{\tilde{S}_j(Q)} \left| t^{2k} L e^{-t^{2k}L} (\chi_{\mathbb{R}^n \setminus 2^{j+j_0+2}Q} m)(y) \right|^2 \frac{dy dt}{t} \\ &=: B_{j,1} + B_{j,2} + B_{j,3}. \end{aligned}$$

From the k -Davies-Gaffney estimate, (4.5) and choosing $\alpha \in (2n(1/p - 1/2)/(1 - \theta), \infty)$, we infer that

$$\begin{aligned} B_{j,1} + B_{j,3} &\lesssim \int_0^{2^{\theta(j-5)l(Q)}} \exp \left\{ -\tilde{C} \left[\frac{2^j l(Q)}{t} \right]^{2k/(2k-1)} \right\} \|m\|_{L^2(\mathbb{R}^n)}^2 \frac{dt}{t} \\ &\lesssim \|m\|_{L^2(\mathbb{R}^n)}^2 \int_0^{2^{\theta(j-5)l(Q)}} \left[\frac{t}{2^j l(Q)} \right]^\alpha \frac{dt}{t} \sim [l(Q)]^{2n(\frac{1}{2}-\frac{1}{p})} [2^{\theta(j-5)-j}]^\alpha, \end{aligned}$$

where \tilde{C} denotes a positive constant. Let $\alpha_1 := (1 - \theta)\alpha/2 - n(1/p - 1/2)$. Then $\alpha_1 \in (0, \infty)$ and we have

$$B_{j,1} + B_{j,3} \lesssim [2^j l(Q)]^{2n(1/2-1/p)} 2^{-2j\alpha_1}. \quad (4.7)$$

Finally, by (4.3) and the size condition (4.2) of m , we conclude that

$$B_{j,2} \lesssim \|m\|_{L^2(\hat{S}_j(Q))}^2 \lesssim \sum_{\ell=j-j_0-2}^{j+j_0+2} \|m\|_{L^2(S_\ell(Q))}^2 \lesssim 2^{-2j\epsilon} [2^j l(Q)]^{2n(1/2-1/p)},$$

which, together with (4.6) and (4.7), shows that there exists a positive constant $\alpha_2 := \min\{\alpha_0, \alpha_1, \epsilon\}$ such that for all $j \in \{j_0 + 5, \dots\}$,

$$I_j \lesssim [2^j l(Q)]^{n(1/2-1/p)} 2^{-j\alpha_2} \sim |S_j(Q)|^{1/2-1/p} 2^{-j\alpha_2}. \quad (4.8)$$

Combining (4.5) and (4.8), we see that

$$\|S_L m\|_{L^p(\mathbb{R}^n)} \lesssim [l(Q)]^{n(\frac{1}{2}-\frac{1}{p})} |2^{j_0+4}Q|^{\frac{1}{p}-\frac{1}{2}} + \sum_{j=j_0+5}^{\infty} 2^{-j\alpha_2} \lesssim 1,$$

from which we deduce (4.4). Thus, the inclusion $\mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n) \subset \mathbb{H}_L^p(\mathbb{R}^n)$ holds, which completes the proof of part one of Theorem 4.5. \square

Now, we prove the inclusion $\mathbb{H}_L^p(\mathbb{R}^n) \subset \mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n)$. To this end, we need to use some results concerning the tent space from [17]. Let F be a function on $\mathbb{R}_+^{n+1} := \mathbb{R}^n \times (0, \infty)$. The \mathcal{A} -functional of F is defined by setting, for all $x \in \mathbb{R}^n$,

$$\mathcal{A}(F)(x) := \left\{ \iint_{\Gamma(x)} |F(y, t)|^2 \frac{dy dt}{t^{n+1}} \right\}^{\frac{1}{2}}.$$

For $p \in (0, \infty)$, the *tent space* $T^p(\mathbb{R}_+^{n+1})$ is defined by

$$T^p(\mathbb{R}_+^{n+1}) := \left\{ F : \mathbb{R}_+^{n+1} \rightarrow \mathbb{C} : \|F\|_{T^p(\mathbb{R}_+^{n+1})} := \|\mathcal{A}(F)\|_{L^p(\mathbb{R}^n)} < \infty \right\}.$$

For any cube Q , denote by $R_Q := Q \times (0, l(Q))$ the *Carleson box* of Q . A measurable function A on \mathbb{R}_+^{n+1} is called a $T^p(\mathbb{R}_+^{n+1})$ -atom associated with Q with $p \in (0, 1]$, if A satisfies the following properties:

$$\text{supp } A \subset R_Q \quad (4.9)$$

and

$$\left\{ \iint_{R_Q} |A(x, t)|^2 \frac{dx dt}{t} \right\}^{1/2} \leq |Q|^{\frac{1}{2} - \frac{1}{p}}. \quad (4.10)$$

For the tent space $T^p(\mathbb{R}_+^{n+1})$ with $p \in (0, 1]$, we have the following atomic decomposition from [17] (see also [39, Proposition 3.25]).

Theorem 4.7 ([17]). *Let $p \in (0, 1]$. For all $F \in T^p(\mathbb{R}_+^{n+1})$, there exist a numerical sequence $\{\lambda_j\}_{j=0}^\infty$ and a sequence $\{A_j\}_{j=0}^\infty$ of $T^p(\mathbb{R}_+^{n+1})$ -atoms such that for almost every $(x, t) \in \mathbb{R}_+^{n+1}$,*

$$F(x, t) = \sum_{j=0}^\infty \lambda_j A_j(x, t).$$

Moreover,

$$\sum_{j=0}^\infty |\lambda_j|^p \sim \|F\|_{T^p(\mathbb{R}_+^{n+1})}^p,$$

where the implicit equivalent positive constants depend only on n . Finally, if $F \in T^p(\mathbb{R}_+^{n+1}) \cap T^2(\mathbb{R}_+^{n+1})$, then the decomposition also converges in $T^2(\mathbb{R}_+^{n+1})$.

Let $M \in \mathbb{N}$. For all $F \in T^2(\mathbb{R}_+^{n+1})$, define the operator $\pi_{M,L}$ by setting, for all $x \in \mathbb{R}^n$,

$$\pi_{M,L}F(x) := \int_0^\infty (t^{2k}L)^{M+1} e^{-t^{2k}L} F(x, t) \frac{dt}{t}. \quad (4.11)$$

For this operator, we have the following useful properties.

Lemma 4.8. *Let $M \in \mathbb{N}$, $p \in (0, 1]$, $\epsilon \in (0, \infty)$ and the operator L satisfy the assumptions (A_1) , (A_2) and (A_3) in Section 2. Let $\pi_{M,L}$ be as in (4.11). Then*

- (i) *The operator $\pi_{M,L}$ is bounded from $T^2(\mathbb{R}_+^{n+1})$ to $L^2(\mathbb{R}^n)$.*
- (ii) *For any $T^p(\mathbb{R}_+^{n+1})$ -atom A , $\pi_{M,L}A$ is an (H_L^p, ϵ, M) -molecule up to a harmless positive constant multiple.*
- (iii) *If $M \in (n(1/p - 1/2)/(2k), \infty)$, then the operator $\pi_{M,L}$ is bounded from the tent space $T^p(\mathbb{R}_+^{n+1})$ to the molecular Hardy space $H_{L, \text{mol}, M}^p(\mathbb{R}^n)$.*

Proof. We first show (i). Let L^* be the adjoint operator of L in $L^2(\mathbb{R}^n)$. Observe that L^* also satisfies the assumptions (A_1) , (A_2) and (A_3) in Section 2. By Fubini's theorem, Hölder's inequality and the quadratic estimate (4.3) with L replaced by L^* , we see that for all $F \in T^2(\mathbb{R}_+^{n+1})$ and $g \in L^2(\mathbb{R}^n)$,

$$\begin{aligned} |(\pi_{M,L}F, g)| &= \left| \int_{\mathbb{R}^n} \int_0^\infty (t^{2k}L)^{M+1} e^{-t^{2k}L} F(x, t) \overline{g(x)} \frac{dt}{t} dx \right| \\ &= \left| \int_0^\infty \int_{\mathbb{R}^n} F(x, t) \overline{(t^{2k}L^*)^{M+1} e^{-t^{2k}L^*} g(x)} dx \frac{dt}{t} \right| \end{aligned}$$

$$\begin{aligned} &\lesssim \left\{ \int_0^\infty \int_{\mathbb{R}^n} |F(x, t)|^2 dx \frac{dt}{t} \right\}^{1/2} \left\{ \int_0^\infty \int_{\mathbb{R}^n} \left| (t^{2k} L^*)^{M+1} e^{-t^{2k} L^*} g(x) \right|^2 dx \frac{dt}{t} \right\}^{1/2} \\ &\lesssim \|F\|_{T^2(\mathbb{R}_+^{n+1})} \|g\|_{L^2(\mathbb{R}^n)}, \end{aligned}$$

which further implies that the operator $\pi_{M,L}$ is bounded from $T^2(\mathbb{R}_+^{n+1})$ to $L^2(\mathbb{R}^n)$. Thus, (i) holds.

To prove (ii), let A be a $T^p(\mathbb{R}_+^{n+1})$ -atom A associated with the cube Q . From (4.11), it follows that for all $\ell \in \{0, \dots, M\}$ and $x \in \mathbb{R}^n$,

$$\pi_{M,L} A(x) = \int_0^\infty (t^{2k} L)^{M+1} e^{-t^{2k} L} A(x, t) \frac{dt}{t} = L^\ell \int_0^\infty t^{2k(M+1)} L^{M+1-\ell} e^{-t^{2k} L} A(x, t) \frac{dt}{t}. \quad (4.12)$$

Observe that

$$\int_0^\infty t^{2k(M+1)} L^{M+1-\ell} e^{-t^{2k} L} A(x, t) \frac{dt}{t} = \int_0^\infty t^{2k(M+1)} \left(L^{1-\frac{\ell}{M+1}} \right)^{M+1} e^{-t^{2k} L} A(x, t) \frac{dt}{t},$$

which belongs to $L^2(\mathbb{R}^n)$ via a dual argument similar to that used in the proof of (i). This, combined with (4.12), implies that $\pi_{M,L}(A)$ satisfies Definition 4.3(i).

For all $x \in \mathbb{R}^n$, letting

$$b(x) := \int_0^\infty t^{2k(M+1)} L e^{-t^{2k} L} A(x, t) \frac{dt}{t},$$

we then have $\pi_{M,L} A(x) = L^M b(x)$. For all $g \in L^2(\mathbb{R}^n)$, from Hölder's inequality, (4.9), (4.10) and the quadratic estimate (4.3) with L replaced by L^* , we deduce that

$$\begin{aligned} &\left| \int_{\mathbb{R}^n} ([l(Q)]^{2k} L)^\ell b(x) \overline{g(x)} dx \right| \\ &= \left| \int_{\mathbb{R}^n} \left\{ \int_0^\infty [l(Q)]^{2k\ell} L^{\ell+1} t^{2k(M+1)} e^{-t^{2k} L} A(x, t) \frac{dt}{t} \right\} \overline{g(x)} dx \right| \\ &= [l(Q)]^{2k\ell} \left| \iint_{R_Q} A(x, t) \overline{(L^*)^{\ell+1} t^{2k(M+1)} e^{-t^{2k} L^*} g(x)} \frac{dx dt}{t} \right| \\ &\lesssim [l(Q)]^{2k\ell} \left[\iint_{R_Q} |A(x, t)|^2 \frac{dx dt}{t} \right]^{1/2} \left[\iint_{R_Q} \left| \overline{(L^*)^{\ell+1} t^{2k(M+1)} e^{-t^{2k} L^*} g(x)} \right|^2 \frac{dx dt}{t} \right]^{1/2} \\ &\lesssim [l(Q)]^{n(1/2-1/p)+2kM} \left\{ \iint_{R_Q} \left| (t^{2k} L^*)^{\ell+1} e^{-t^{2k} L^*} g(x) \right|^2 \frac{dx dt}{t} \right\}^{1/2} \\ &\lesssim [l(Q)]^{n(1/2-1/p)+2kM} \|g\|_{L^2(\mathbb{R}^n)}, \end{aligned} \quad (4.13)$$

which further implies that for all $\ell \in \{0, \dots, M\}$,

$$\left\| ([l(Q)]^{2k} L)^\ell b \right\|_{L^2(2Q)} \lesssim [l(Q)]^{2kM} |Q|^{\frac{1}{2}-\frac{1}{p}}.$$

Thus, by this, we conclude that for all $\tilde{\ell} \in \{0, \dots, M\}$ and $j \in \{0, 1\}$,

$$\begin{aligned} \left\| ([l(Q)]^{-2k} L^{-1})^{\tilde{\ell}} (\pi_{M,L} A) \right\|_{L^2(S_j(Q))} &= \left\| ([l(Q)]^{-2k} L^{-1})^{\tilde{\ell}} L^M b \right\|_{L^2(S_j(Q))} \\ &\lesssim \left\| ([l(Q)]^{2k(M-\tilde{\ell})} L^{M-\tilde{\ell}}) b \right\|_{L^2(2(Q))} |l(Q)|^{-2kM} \\ &\lesssim [l(Q)]^{n(1/2-1/p)}, \end{aligned} \quad (4.14)$$

which is desired.

Moreover, for all $\tilde{\ell} \in \{0, \dots, M\}$ and $j \in \{2, 3, \dots\}$, letting $g \in L^2(\mathbb{R}^n)$ with $\text{supp } g \subset S_j(Q)$, choosing $\alpha \in (n(1/p - 1/2)(2 - 1/k), \infty)$ and using Lemma 3.1, similar to the estimate for (4.13), we see that

$$\begin{aligned} & \left| \int_{\mathbb{R}^n} ([l(Q)]^{2k} L)^\ell b(x) \overline{g(x)} dx \right| \\ & \lesssim [l(Q)]^{n(1/2-1/p)+2kM} \left\{ \int_0^{l(Q)} \int_Q \left| (t^{2k} L^*)^{\ell+1} e^{-t^{2k} L^*} g(x) \right|^2 \frac{dx dt}{t} \right\}^{1/2} \\ & \lesssim [l(Q)]^{n(1/2-1/p)+2kM} \left[\int_0^{l(Q)} \exp \left\{ -C \left[\frac{\text{dist}(Q, S_j(Q))}{t} \right]^{2k/(2k-1)} \right\} \frac{dt}{t} \right]^{1/2} \|g\|_{L^2(S_j(Q))} \\ & \lesssim [l(Q)]^{n(1/2-1/p)+2kM} 2^{-jk\alpha/(2k-1)} \|g\|_{L^2(S_j(Q))}, \end{aligned}$$

which further implies that for all $\tilde{\ell} \in \{0, \dots, M\}$ and $j \in \{2, 3, \dots\}$,

$$\begin{aligned} \left\| ([l(Q)]^{-2k} L^{-1})^{\tilde{\ell}} (\pi_{M,L} A) \right\|_{L^2(S_j(Q))} & \leq \left\| ([l(Q)]^{-2k} L^{-1})^{\tilde{\ell}} L^M b \right\|_{L^2(S_j(Q))} \\ & \lesssim \left\| ([l(Q)]^{2k(M-\tilde{\ell})} L^{M-\tilde{\ell}}) b \right\|_{L^2(S_j(Q))} |l(Q)|^{-2kM} \\ & \lesssim [2^j l(Q)]^{n(1/2-1/p)} 2^{-j[k\alpha/(2k-1)-n(1/p-1/2)]} \\ & \sim [2^j l(Q)]^{n(1/2-1/p)} 2^{-j\epsilon}, \end{aligned} \quad (4.15)$$

where $\epsilon := k\alpha/(2k-1) - n(1/p - 1/2) \in (0, \infty)$.

Combining (4.14) and (4.15), we know that $\pi_{M,L} A$ satisfies Definition 4.3(ii) up to a harmless positive constant multiple. Thus, $\pi_{M,L} A$ is an (H_L^p, ϵ, M) -molecule up to a harmless positive constant multiple, which completes the proof of (ii).

To show (iii), by density, we only need to show that for all $F \in T^p(\mathbb{R}_+^{n+1}) \cap T^2(\mathbb{R}_+^{n+1})$,

$$\|\pi_{M,L} F\|_{H_{L, \text{mol}, M}^p(\mathbb{R}^n)} \lesssim \|F\|_{T^p(\mathbb{R}_+^{n+1})}.$$

To this end, by Theorem 4.7, there exist a sequence $\{A_i\}_{i=0}^\infty$ of $T^p(\mathbb{R}_+^{n+1})$ -atoms and $\{\lambda_i\}_{i=0}^\infty \in l^p$ such that $F = \sum_{i=0}^\infty \lambda_i A_i$ in both pointwise and $T^2(\mathbb{R}_+^{n+1})$, and

$$\left(\sum_{i=0}^\infty |\lambda_i|^p \right)^{1/p} \sim \|F\|_{T^p(\mathbb{R}_+^{n+1})}.$$

By (i) of this lemma, we know that

$$\pi_{M,L} F = \sum_{i=0}^\infty \lambda_i \pi_{M,L} A_i$$

in $L^2(\mathbb{R}^n)$, which, combining (ii) of Lemma 4.8, shows that $\sum_{i=0}^\infty \lambda_i \pi_{M,L} A_i$ is a molecular $(H_L^p, 2, \epsilon, M)$ -representation of $\pi_{M,L} F$. Thus, $\pi_{M,L} F \in H_{L, \text{mol}, M}^p(\mathbb{R}^n)$ and

$$\|\pi_{M,L} F\|_{H_{L, \text{mol}, M}^p(\mathbb{R}^n)} \lesssim \left\{ \sum_{i=0}^\infty |\lambda_i|^p \right\}^{1/p} \sim \|F\|_{T^p(\mathbb{R}_+^{n+1})},$$

which completes the proof of (iii) and hence Lemma 4.8. \square

Proof of Theorem 4.5: the inclusion $\mathbb{H}_L^p(\mathbb{R}^n) \subset \mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n)$. For all $f \in \mathbb{H}_L^p(\mathbb{R}^n)$, $t \in (0, \infty)$ and $x \in \mathbb{R}^n$, let

$$F(x, t) := t^{2k} L e^{-t^{2k} L} f(x). \quad (4.16)$$

By $S_L f \in L^p(\mathbb{R}^n)$ and $f \in L^2(\mathbb{R}^n)$, together with the fact that S_L is bounded on $L^2(\mathbb{R}^n)$, we know that $F \in T^p(\mathbb{R}_+^{n+1}) \cap T^2(\mathbb{R}_+^{n+1})$. Moreover, by the H_∞ functional calculus in $L^2(\mathbb{R}^n)$, we see the following Calderón reproducing formula that for all $g \in L^2(\mathbb{R}^n)$,

$$g = C_9 \int_0^\infty (t^{2k} L)^{M+2} e^{-2t^{2k} L} g \frac{dt}{t},$$

where C_9 is a positive constant such that $C_9 \int_0^\infty t^{2k(M+2)} e^{-2t^{2k}} \frac{dt}{t} = 1$. Thus, for all $f \in \mathbb{H}_L^p(\mathbb{R}^n)$, if letting F be as in (4.16), then $f = C_9 \pi_{M,L} F$ and, by Lemma 4.8(iii) and its proof, we further know that $f \in \mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n)$ and $\|f\|_{\mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n)}^p \lesssim \|f\|_{\mathbb{H}_L^p(\mathbb{R}^n)}^p$. Therefore, $\mathbb{H}_L^p(\mathbb{R}^n) \subset \mathbb{H}_{L, \text{mol}, M}^p(\mathbb{R}^n)$, which completes the proof of Theorem 4.5. \square

5 Generalized square function characterizations of $H_L^p(\mathbb{R}^n)$

This section is devoted to the generalized square function characterization of $H_L^p(\mathbb{R}^n)$. We first introduce the notion of the Hardy space $H_{\psi,L}^p(\mathbb{R}^n)$ defined via the generalized square function. Let $\omega \in [0, \pi/2)$, $\alpha \in (0, \infty)$, $\beta \in (n(1/p - 1/2)/(2k), \infty)$ and $\psi \in \Psi_{\alpha,\beta}(S_\mu^0)$ with $\mu \in (\omega, \pi/2)$. For all $f \in L^2(\mathbb{R}^n)$ and $(x, t) \in \mathbb{R}_+^{n+1}$, define the operator $Q_{\psi,L} f$ by,

$$Q_{\psi,L} f(x, t) := \psi(t^{2k} L) f(x). \quad (5.1)$$

Definition 5.1. Let $p \in (0, 1]$, $\omega \in [0, \pi/2)$, L be the operator of type ω satisfying the assumptions (A_1) , (A_2) and (A_3) in Section 2, $\alpha \in (0, \infty)$, $\beta \in (n(1/p - 1/2)/(2k), \infty)$, $\mu \in (\omega, \pi/2)$ and $\psi \in \Psi_{\alpha,\beta}(S_\mu^0)$. The *generalized square function Hardy space* $H_{\psi,L}^p(\mathbb{R}^n)$ is defined to be the completion of the space

$$\mathbb{H}_{\psi,L}^p(\mathbb{R}^n) := \{f \in L^2(\mathbb{R}^n) : Q_{\psi,L} f \in T^p(\mathbb{R}_+^{n+1})\}$$

with respect to the *quasi-norm* $\|f\|_{H_{\psi,L}^p(\mathbb{R}^n)} := \|Q_{\psi,L} f\|_{T^p(\mathbb{R}_+^{n+1})}$.

The following theorem, which establishes the generalized square function characterization of $H_L^p(\mathbb{R}^n)$, is the main result of this section.

Theorem 5.2. Let $p \in (0, 1]$, $\omega \in [0, \pi/2)$, L be the operator of type ω satisfying the assumptions (A_1) , (A_2) and (A_3) in Section 2, $\alpha \in (0, \infty)$, $\beta \in (n(1/p - 1/2)/(2k), \infty)$, $\mu \in (\omega, \pi/2)$ and $\psi \in \Psi_{\alpha,\beta}(S_\mu^0)$. Then the Hardy space $H_L^p(\mathbb{R}^n) = H_{\psi,L}^p(\mathbb{R}^n)$ with equivalent norms.

Before proving Theorem 5.2, we first give an application of this theorem. Let $\alpha \in (0, \infty)$ and L^α be the *fractional power with exponent α of L* defined by the H_∞ functional calculus in $L^2(\mathbb{R}^n)$ (see, for example, [33, 46]). More precisely, choose $m \in \mathbb{N}$ such that $m > \alpha$. Then, $z^\alpha(1+z)^{-m} \in \Psi_{\alpha, m-\alpha}(S_\mu^0)$ for all $\mu \in [0, \pi/2)$ and L^α is defined by setting

$$L^\alpha := (z^\alpha)(L) := (1+L)^m \left(\frac{z^\alpha}{(1+z)^m} \right) (L).$$

For more details about L^α , we refer the reader to [33, 46] and the references therein.

Assume that $-L^\alpha$ generates a bounded holomorphic semigroup $\{e^{-tL^\alpha}\}_{t>0}$. From [33, Example 3.4.6], it follows that this is true when $\alpha \in (0, 1]$, and in this case, $\{e^{-tL^\alpha}\}_{t>0}$ is called the *subordinated semigroup* (see [33, p. 80] for more details). For all $f \in L^2(\mathbb{R}^n)$, define the L^α -adapted square function S_{L^α} by setting, for all $x \in \mathbb{R}^n$,

$$S_{L^\alpha} f(x) := \left\{ \iint_{\Gamma(x)} \left| t^{2k\alpha} L^\alpha e^{-t^{2k\alpha} L^\alpha} f(y) \right|^2 \frac{dy dt}{t^{n+1}} \right\}^{1/2}. \quad (5.2)$$

For $p \in (0, 1]$, we also define the *Hardy space* $H_{L^\alpha}^p(\mathbb{R}^n)$ associated with L^α to be the completion of the set

$$\mathbb{H}_{L^\alpha}^p(\mathbb{R}^n) := \left\{ f \in L^2(\mathbb{R}^n) : \|S_{L^\alpha} f\|_{L^p(\mathbb{R}^n)} < \infty \right\} \quad (5.3)$$

with respect to the quasi-norm $\|f\|_{H_{L^\alpha}^p(\mathbb{R}^n)} := \|S_{L^\alpha} f\|_{L^p(\mathbb{R}^n)}$.

With the help of Theorem 5.2, we immediately obtain the following interesting corollary.

Corollary 5.3. *Let $p \in (0, 1]$ and L satisfy the assumptions (A_1) , (A_2) and (A_3) . Assume further that when $\alpha \in (1, \infty)$, $-L^\alpha$ generates a bounded holomorphic semigroup. Then, for all $\alpha \in (0, \infty)$, the Hardy spaces $H_{L^\alpha}^p(\mathbb{R}^n) = H_L^p(\mathbb{R}^n)$ with equivalent norms.*

Proof. Let $\omega \in [0, \pi/2)$. Recall that L is an operator of type ω . For all $\alpha \in (0, \infty)$, $\mu \in (\omega, \pi/2)$ and $\xi \in S_\mu^0$, set $\psi(\xi) := \xi^\alpha e^{-\xi^\alpha}$. Then, for all $\beta \in (n(1/p - 1/2)/(2k), \infty)$, $\psi \in \Psi_{\alpha, \beta}(S_\mu^0)$ and hence, by Theorem 5.2, we conclude that for all $f \in L^2(\mathbb{R}^n)$,

$$\|f\|_{H_{L^\alpha}^p(\mathbb{R}^n)} = \|S_{L^\alpha} f\|_{L^p(\mathbb{R}^n)} = \|Q_{\psi, L} f\|_{T^p(\mathbb{R}_+^{n+1})} = \|f\|_{H_{\psi, L}^p(\mathbb{R}^n)} \sim \|f\|_{H_L^p(\mathbb{R}^n)},$$

which, together with the density of $L^2(\mathbb{R}^n)$ in $H_L^p(\mathbb{R}^n)$ and $H_{L^\alpha}^p(\mathbb{R}^n)$, shows that $H_L^p(\mathbb{R}^n) = H_{L^\alpha}^p(\mathbb{R}^n)$ with equivalent norms. This finishes the proof of Corollary 5.3. \square

Let $\omega \in [0, \pi/2)$ be as in Section 2 and $\mu \in (\omega, \pi/2)$. To prove Theorem 5.2, we introduce two operators as follows:

(i) For all $F \in T^2(\mathbb{R}_+^{n+1})$ and $\psi \in \Psi(S_\mu^0)$, the operator $\pi_{\psi, L}$ is defined by setting, for all $x \in \mathbb{R}^n$,

$$\pi_{\psi, L} F(x) := \int_0^\infty \psi(t^{2k} L) F(x, t) \frac{dt}{t}; \quad (5.4)$$

(ii) For all $\psi, \tilde{\psi} \in \Psi(S_\mu^0)$, $f \in H_\infty(S_\mu^0)$ and $F \in T^2(\mathbb{R}_+^{n+1})$, the operator Q^f is defined by setting, for all $x \in \mathbb{R}^n$ and $s \in (0, \infty)$,

$$Q^f F(x, s) := Q_{\psi, L} \circ f(L) \circ \pi_{\tilde{\psi}, L} F(x, s) = \int_0^\infty \psi(s^{2k} L) f(L) \tilde{\psi}(t^{2k} L) F(x, t) \frac{dt}{t}, \quad (5.5)$$

where the operator $Q_{\psi, L}$ is defined as in (5.1).

Observe that by (4.3), $Q_{\psi, L}$ is bounded from $L^2(\mathbb{R}^n)$ to $T^2(\mathbb{R}_+^{n+1})$ and so is $Q_{\tilde{\psi}, L^*}$. By Fubini's theorem and Hölder's inequality, we see that for all $F \in T^2(\mathbb{R}_+^{n+1})$ and $g \in L^2(\mathbb{R}^n)$,

$$\int_{\mathbb{R}^n} \pi_{\psi, L} F(x) \overline{g(x)} dx = \int_{\mathbb{R}^n} \int_0^\infty \psi(t^{2k} L) F(x, t) \frac{dt}{t} \overline{g(x)} dx = \int_0^\infty \int_{\mathbb{R}^n} F(x, t) \overline{Q_{\tilde{\psi}, L^*}(g)(x)} dx \frac{dt}{t}.$$

Thus, $Q_{\tilde{\psi}, L^*}$ is the adjoint operator of $\pi_{\psi, L}$, which, together with the above observation, shows that $\pi_{\psi, L}$ is bounded from $T^2(\mathbb{R}_+^{n+1})$ to $L^2(\mathbb{R}^n)$. From these facts and (5.5) together with that L has a bounded H_∞ functional calculus in $L^2(\mathbb{R}^n)$, it follows that Q^f is bounded on $T^2(\mathbb{R}_+^{n+1})$.

Let $\sigma_1, \sigma_2, \tau_1, \tau_2 \in (0, \infty)$. Assume that $\psi \in \Psi_{\sigma_1, \tau_1}(S_\mu^0)$ and $\tilde{\psi} \in \Psi_{\sigma_2, \tau_2}(S_\mu^0)$. We now consider the operator $\psi(s^{2k} L) f(L) \tilde{\psi}(t^{2k} L)$ in (5.5). Let $a \in (0, \min\{\sigma_1, \tau_2\})$ and $b \in (0, \min\{\sigma_2, \tau_1\})$. For $s, t \in (0, \infty)$, when $s \leq t$, we write

$$\psi(s^{2k} L) f(L) \tilde{\psi}(t^{2k} L) = \left(\frac{s^{2k}}{t^{2k}} \right)^a (s^{2k} L)^{-a} \psi(s^{2k} L) f(L) (t^{2k} L)^a \tilde{\psi}(t^{2k} L) =: \left(\frac{s^{2k}}{t^{2k}} \right)^a T_{s^{2k}, t^{2k}}, \quad (5.6)$$

while when $s > t$, we write

$$\psi(s^{2k} L) f(L) \tilde{\psi}(t^{2k} L) = \left(\frac{t^{2k}}{s^{2k}} \right)^b (s^{2k} L)^b \psi(s^{2k} L) f(L) (t^{2k} L)^{-b} \tilde{\psi}(t^{2k} L) =: \left(\frac{t^{2k}}{s^{2k}} \right)^b T_{s^{2k}, t^{2k}}. \quad (5.7)$$

Then, we have the following useful estimate on $\{T_{s, t}\}_{s, t > 0}$.

Lemma 5.4. Let $k \in \mathbb{N}$ be as in (2.6), $\sigma_1, \sigma_2, \tau_1, \tau_2 \in (0, \infty)$, $\omega \in [0, \pi/2)$, $\mu \in (\omega, \pi/2)$, $\psi \in \Psi_{\sigma_1, \tau_1}(S_\mu^0)$, $\tilde{\psi} \in \Psi_{\sigma_2, \tau_2}(S_\mu^0)$, $a \in (0, \min\{\sigma_1, \tau_2\})$ and $b \in (0, \min\{\sigma_2, \tau_1\})$. Let $f \in H_\infty(S_\mu^0)$. Let $\{T_{s,t}\}_{s,t>0}$ be as in (5.6) and (5.7) with s^{2k} and t^{2k} replaced, respectively, by s and t . Then, there exists a positive constant C such that for all $M \in (0, \min\{\sigma_2 + a, \tau_1 + b\})$, $s, t \in (0, \infty)$, closed sets $E, F \subset \mathbb{R}^n$ and $g \in L^2(\mathbb{R}^n)$ supported in E ,

$$\|T_{s,t}g\|_{L^2(F)} \leq C\|f\|_{L^\infty(\mathbb{R}^n)} \min \left\{ 1, \frac{\max\{t, s\}}{[\text{dist}(E, F)]^{2k}} \right\}^M \|g\|_{L^2(E)}. \quad (5.8)$$

Proof. We prove this lemma by considering two cases. If $s \leq t$, since $a \in (0, \min\{\sigma_1, \tau_2\})$, we conclude that for all $\xi \in S_\mu^0$,

$$\left| (s\xi)^{-a} \psi(s\xi) f(\xi) \right| \lesssim \frac{|s\xi|^{\sigma_1-a}}{1 + |s\xi|^{\sigma_1+\tau_1}} \|f\|_{L^\infty(\mathbb{R}^n)} \lesssim 1$$

and

$$\left| (t\xi)^a \tilde{\psi}(t\xi) \right| \lesssim \frac{|t\xi|^{\sigma_2+a}}{1 + |t\xi|^{\sigma_2+\tau_2}},$$

which, together with Lemma 3.3 with ψ and f therein replaced by $(t\xi)^a \tilde{\psi}(t\xi)$ and $(s\xi)^{-a} \psi(s\xi) f(\xi)$ respectively, implies that the family $\{T_{s,t}\}_{s \leq t}$ of operators satisfies the k -Davies-Gaffney estimate of order $\sigma_2 + a$ in t .

Similarly, if $s > t$, since $b \in (0, \min\{\sigma_2, \tau_1\})$, we see that for all $\xi \in S_\mu^0$,

$$\left| f(\xi) (t\xi)^{-b} \tilde{\psi}(t\xi) \right| \leq \frac{|t\xi|^{\sigma_2-b}}{1 + |t\xi|^{\sigma_2+\tau_2}} \|f\|_{L^\infty(\mathbb{R}^n)} \lesssim 1$$

and

$$\left| (s\xi)^b \psi(s\xi) \right| \leq \frac{|s\xi|^{\tau_1+b}}{1 + |s\xi|^{\sigma_1+\tau_1}},$$

which, together with Lemma 3.3 with ψ and f therein replaced by $(s\xi)^b \psi(s\xi)$ and $f(\xi)(t\xi)^{-b} \tilde{\psi}(t\xi)$ respectively, implies that the family $\{T_{s,t}\}_{s > t}$ of operators satisfies the k -Davies-Gaffney estimate of order $\tau_1 + b$ in s .

Thus, for all $M \in (0, \min\{\sigma_2 + a, \tau_1 + b\})$, we immediately obtain (5.8), which completes the proof of Lemma 5.4. \square

Lemma 5.5. Let $p \in (0, 1]$, L be the operator of type ω satisfying the assumptions (A_1) , (A_2) and (A_3) in Section 2, $\alpha \in (0, \infty)$, $\beta \in (n(1/p - 1/2)/(2k), \infty)$, $\omega \in [0, \pi/2)$, $\mu \in (\omega, \pi/2)$, $\psi \in \Psi_{\alpha, \beta}(S_\mu^0)$ and $\tilde{\psi} \in \Psi_{\beta, \alpha}(S_\mu^0)$. Then the operator Q^f originally defined in (5.5) on $T^2(\mathbb{R}_+^{n+1})$ can be continuously extended to a bounded linear operator on $T^p(\mathbb{R}_+^{n+1})$. Moreover, there exists a positive constant C such that for all $F \in T^p(\mathbb{R}_+^{n+1})$ and $f \in H_\infty(S_\mu^0)$,

$$\|Q^f F\|_{T^p(\mathbb{R}_+^{n+1})} \leq C\|f\|_{L^\infty(S_\mu^0)} \|F\|_{T^p(\mathbb{R}_+^{n+1})}. \quad (5.9)$$

Proof. By the density of $T^2(\mathbb{R}_+^{n+1}) \cap T^p(\mathbb{R}_+^{n+1})$ in $T^p(\mathbb{R}_+^{n+1})$ (see [17]), it suffices to prove (5.9) for all $F \in T^2(\mathbb{R}_+^{n+1}) \cap T^p(\mathbb{R}_+^{n+1})$. To this end, by borrowing some ideas from the proof of Theorem 1.1 in [54], we only need to show that for all $T^p(\mathbb{R}_+^{n+1})$ -atoms A ,

$$\|Q^f A\|_{T^p(\mathbb{R}_+^{n+1})} \lesssim \|f\|_{L^\infty(S_\mu^0)}. \quad (5.10)$$

Indeed, if (5.10) holds, then from Theorem 4.7 and the $T^2(\mathbb{R}_+^{n+1})$ -boundedness of $Q^f A$, it follows that for any $F \in T^2(\mathbb{R}_+^{n+1}) \cap T^p(\mathbb{R}_+^{n+1})$, there exist a sequence $\{A_j\}_{j=0}^\infty$ of $T^p(\mathbb{R}_+^{n+1})$ -atoms and $\{\lambda_j\}_{j=0}^\infty \in$

l^p such that $F := \sum_{j=0}^{\infty} \lambda_j A_j$ with the summation converges in both pointwise and $T^2(\mathbb{R}_+^{n+1})$, and $\{\sum_{j=0}^{\infty} |\lambda_j|^p\}^{1/p} \sim \|F\|_{T^p(\mathbb{R}_+^{n+1})}$. We claim that for $\frac{dx dt}{t}$ -almost every $(x, t) \in \mathbb{R}_+^{n+1}$,

$$\left| Q^f \left(\sum_{j=0}^{\infty} \lambda_j A_j \right) (x, t) \right| \leq \sum_{j=0}^{\infty} |\lambda_j Q^f A_j(x, t)|. \quad (5.11)$$

Assume this claim for the moment. By (5.11) and $p \in (0, 1]$, together with the monotonicity of l^p , we have

$$\begin{aligned} \|Q^f F\|_{T^p(\mathbb{R}_+^{n+1})} &\leq \left\{ \sum_{j=0}^{\infty} |\lambda_j|^p \|Q^f A_j\|_{T^p(\mathbb{R}_+^{n+1})}^p \right\}^{1/p} \\ &\leq \sup_{j \in \mathbb{Z}_+} \left\{ \|Q^f(A_j)\|_{T^p(\mathbb{R}_+^{n+1})} \right\} \left\{ \sum_{j=0}^{\infty} |\lambda_j|^p \right\}^{1/p} \\ &\lesssim \|f\|_{L^\infty(S_\mu^0)} \left\{ \sum_{j=0}^{\infty} |\lambda_j|^p \right\}^{1/p} \sim \|f\|_{L^\infty(S_\mu^0)} \|F\|_{T^p(\mathbb{R}_+^{n+1})}. \end{aligned}$$

That is, Q^f is bounded on $T^p(\mathbb{R}_+^{n+1})$. To show the claim (5.11), for simplicity of the notation, let $d\mu(x, t) := \frac{dx dt}{t}$ for all $(x, t) \in \mathbb{R}_+^{n+1}$. By the $T^2(\mathbb{R}_+^{n+1})$ -boundedness of Q^f and the $T^2(\mathbb{R}_+^{n+1})$ -convergence of $F = \sum_{j=0}^{\infty} \lambda_j A_j$, we conclude that for any $\eta \in (0, \infty)$,

$$\lim_{N \rightarrow \infty} \mu \left(\left\{ x \in \mathbb{R}^n : \left| Q^f \left(\sum_{i=N+1}^{\infty} \lambda_i A_i \right) \right| > \eta \right\} \right) \lesssim \lim_{N \rightarrow \infty} \frac{1}{\eta^2} \left\| \sum_{i=N+1}^{\infty} \lambda_i A_i \right\|_{T^2(\mathbb{R}_+^{n+1})}^2 = 0.$$

This, combined with the Riesz theorem, implies that there exists a subsequence

$$\left\{ Q^f \left(\sum_{j=N_\ell+1}^{\infty} \lambda_j A_j \right) \right\}_{\ell \in \mathbb{N}}$$

of $\{Q^f(\sum_{j=N+1}^{\infty} \lambda_j A_j)\}_{N \in \mathbb{N}}$ such that for μ -almost every $(x, t) \in \mathbb{R}_+^{n+1}$,

$$\lim_{\ell \rightarrow \infty} Q^f \left(\sum_{j=N_\ell+1}^{\infty} \lambda_j A_j \right) (x, t) = 0,$$

where $\{N_\ell\}_{\ell \in \mathbb{N}} \subset \mathbb{N}$ and $\lim_{\ell \rightarrow \infty} N_\ell = \infty$. Therefore, for μ -almost every $(x, t) \in \mathbb{R}_+^{n+1}$ and all $\ell \in \mathbb{N}$,

$$\left| Q^f \left(\sum_{j=0}^{\infty} \lambda_j A_j \right) (x, t) \right| \leq \sum_{j=0}^{N_\ell} |\lambda_j Q^f A_j(x, t)| + \left| Q^f \left(\sum_{j=N_\ell+1}^{\infty} \lambda_j A_j \right) (x, t) \right|,$$

which, together with letting $\ell \rightarrow \infty$, shows the claim (5.11).

To finish the proof of Lemma 5.5, we still need to prove (5.10). By the homogeneity of the norm $\|\cdot\|_{T^p(\mathbb{R}_+^{n+1})}$, without loss of generality, we may assume that $\|f\|_{L^\infty(S_\mu^0)} = 1$. Let Q be the cube associated with the $T^p(\mathbb{R}_+^{n+1})$ -atom A and $R_Q := Q \times (0, l(Q))$, where $l(Q)$ denotes the *side length* of Q . For all $i \in \mathbb{N}$, set $2^i R_Q := 2^i Q \times (0, 2^i l(Q)) \subset \mathbb{R}_+^{n+1}$ and $S_i(R_Q) := 2^i R_Q \setminus (2^{i-1} R_Q)$.

For $i = 1$, by Hölder's inequality, the $T^2(\mathbb{R}_+^{n+1})$ -boundedness of Q^f and the size condition (4.10) of $T^p(\mathbb{R}_+^{n+1})$ -atoms, we see that

$$\|\chi_{2R_Q} Q^f A\|_{T^p(\mathbb{R}_+^{n+1})} = \|\mathcal{A}(\chi_{2R_Q} Q^f A)\|_{L^p(\mathbb{R}^n)}$$

$$\begin{aligned}
&\leq \|\mathcal{A}(\chi_{2R_Q} Q^f A)\|_{L^2(\mathbb{R}^n)} |2(\sqrt{n}+2)Q|^{1/p-1/2} \\
&\lesssim \left\{ \iint_{R_Q} |A(x, t)|^2 \frac{dx dt}{t} \right\}^{1/2} |Q|^{1/p-1/2} \lesssim 1.
\end{aligned} \tag{5.12}$$

For $i \geq 2$, using Hölder's inequality and Fubini's theorem, we then conclude that

$$\begin{aligned}
\|\chi_{S_i(R_Q)} Q^f A\|_{T^p(\mathbb{R}_+^{n+1})} &= \|\mathcal{A}(\chi_{S_i(R_Q)} Q^f A)\|_{L^p(\mathbb{R}^n)} \lesssim \|\mathcal{A}(\chi_{S_i(R_Q)} Q^f A)\|_{L^2(\mathbb{R}^n)} |2^i(2+\sqrt{n})Q|^{\frac{1}{p}-\frac{1}{2}} \\
&\sim \left\{ \left[\int_0^{2^{i-1}l(Q)} \int_{\mathbb{R}^n} \chi_{S_i(R_Q)}(x, s) |Q^f A(x, s)|^2 \frac{dx ds}{s} \right]^{1/2} \right. \\
&\quad \left. + \left[\int_{2^{i-1}l(Q)}^{2^i l(Q)} \int_{\mathbb{R}^n} \cdots \right]^{1/2} \right\} |2^i Q|^{\frac{1}{p}-\frac{1}{2}} \\
&=: \{I + O\} |2^i Q|^{\frac{1}{p}-\frac{1}{2}}.
\end{aligned}$$

To estimate O , from (5.5), Minkowski's inequality, Fubini's inequality, Lemma 5.4 and Hölder's inequality, we deduce that

$$\begin{aligned}
O &\sim \left\{ \int_{2^{i-1}l(Q)}^{2^i l(Q)} \int_{\mathbb{R}^n} \chi_{S_i(R_Q)}(x, s) |Q^f A(x, s)|^2 dx \frac{ds}{s} \right\}^{1/2} \\
&\sim \left\{ \int_{2^{i-1}l(Q)}^{2^i l(Q)} \int_{\mathbb{R}^n} \chi_{S_j(R_Q)}(x, s) \left| \int_0^\infty \psi(s^{2k}L) f(L) \tilde{\psi}(t^{2k}L) A(x, t) \frac{dt}{t} \right|^2 dx \frac{ds}{s} \right\}^{1/2} \\
&\sim \left\{ \int_{2^{i-1}l(Q)}^{2^i l(Q)} \int_{\mathbb{R}^n} \chi_{S_i(R_Q)}(x, s) \left| \int_0^\infty \left(\frac{t}{s}\right)^{2kb} T_{s^{2k}, t^{2k}} A(x, t) \frac{dt}{t} \right|^2 dx \frac{ds}{s} \right\}^{1/2} \\
&\lesssim \int_0^\infty \left[\frac{t}{2^i l(Q)} \right]^{2kb} \left[\int_{2^{i-1}l(Q)}^{2^i l(Q)} \int_{\mathbb{R}^n} |T_{s^{2k}, t^{2k}} A(x, t)|^2 \chi_{S_i(R_Q)}(x, s) dx \frac{ds}{s} \right]^{1/2} \frac{dt}{t} \\
&\lesssim \int_0^{l(Q)} \left[\frac{t}{2^i l(Q)} \right]^{2kb} \|A(\cdot, t)\|_{L^2(\mathbb{R}^n)} \frac{dt}{t} \\
&\lesssim \left\{ \int_0^{l(Q)} \|A(\cdot, t)\|_{L^2(\mathbb{R}^n)}^2 \frac{dt}{t} \right\}^{1/2} \left\{ \int_0^{l(Q)} \left[\frac{t}{2^i l(Q)} \right]^{4kb} \frac{dt}{t} \right\}^{1/2} \\
&\lesssim 2^{-2ikb} |Q|^{1/2-1/p} \sim 2^{-i[2kb-n(1/p-1/2)]} |2^i Q|^{1/2-1/p} \sim 2^{-i\gamma_1} |2^i Q|^{1/2-1/p},
\end{aligned}$$

where $b \in (n(1/p-1/2)/(2k), \beta)$ and $\gamma_1 := 2kb - n(1/p-1/2) > 0$.

Let $a \in (0, \alpha)$. To estimate I , by Fubini's theorem and Minkowski's inequality, we see that

$$\begin{aligned}
I &\sim \left\{ \int_0^{2^{i-1}l(Q)} \int_{\mathbb{R}^n} \chi_{S_i(R_Q)}(x, s) |Q^f A(x, s)|^2 dx \frac{ds}{s} \right\}^{1/2} \\
&\sim \left\{ \int_0^{2^{i-1}l(Q)} \int_{\mathbb{R}^n} \chi_{S_i(R_Q)}(x, s) \left| \int_0^\infty \psi(s^{2k}L) f(L) \tilde{\psi}(t^{2k}L) A(x, t) \frac{dt}{t} \right|^2 dx \frac{ds}{s} \right\}^{1/2} \\
&\sim \left\{ \int_0^{2^{i-1}l(Q)} \int_{\mathbb{R}^n} \chi_{S_i(R_Q)}(x, s) \left| \int_0^\infty \min \left\{ \left(\frac{s}{t}\right)^{2ka}, \left(\frac{t}{s}\right)^{2kb} \right\} T_{s^{2k}, t^{2k}} A(x, t) \frac{dt}{t} \right|^2 dx \frac{ds}{s} \right\}^{1/2} \\
&\lesssim \int_0^{l(Q)} \left\{ \int_0^t \int_{\mathbb{R}^n} \left(\frac{s}{t}\right)^{4ka} |T_{s^{2k}, t^{2k}} A(x, t)|^2 \chi_{S_i(R_Q)}(x, s) dx \frac{ds}{s} \right\}^{1/2} \frac{dt}{t}
\end{aligned}$$

$$+ \int_0^{l(Q)} \left\{ \int_t^{2^{i-1}l(Q)} \int_{\mathbb{R}^n} \left(\frac{t}{s} \right)^{4kb} |T_{s^{2k}, t^{2k}} A(x, t)|^2 \chi_{S_i(R_Q)}(x, s) dx \frac{ds}{s} \right\}^{1/2} \frac{dt}{t}$$

$$=: I_1 + I_2.$$

Let $M \in (n(1/p - 1/2)/(2k), \min\{\alpha + b, \beta + a\})$. It follows, from Lemma 5.4 and Hölder's inequality, that

$$\begin{aligned} I_1 &\lesssim \int_0^{l(Q)} \left[\int_0^t \left(\frac{s}{t} \right)^{4ka} \left\{ \frac{t^{2k}}{[\text{dist}(R_Q, S_i(R_Q))]^{2k}} \right\}^{2M} \|A(\cdot, t)\|_{L^2(\mathbb{R}^n)}^2 \frac{ds}{s} \right]^{1/2} \frac{dt}{t} \\ &\lesssim \int_0^{l(Q)} \left[\int_0^t \left(\frac{s}{t} \right)^{4ka} \left\{ \frac{t^{2k}}{[2^i l(Q)]^{2k}} \right\}^{2M} \|A(\cdot, t)\|_{L^2(\mathbb{R}^n)}^2 \frac{ds}{s} \right]^{1/2} \frac{dt}{t} \\ &\sim \frac{1}{[2^i l(Q)]^{2kM}} \int_0^{l(Q)} \|A(\cdot, t)\|_{L^2(\mathbb{R}^n)} t^{2kM} \left[\int_0^t \left(\frac{s}{t} \right)^{4ka} \frac{ds}{s} \right]^{1/2} \frac{dt}{t} \\ &\lesssim \frac{1}{[2^i l(Q)]^{2kM}} \left\{ \iint_{R_Q} |A(x, t)|^2 \frac{dx dt}{t} \right\}^{1/2} \left\{ \int_0^{l(Q)} t^{4kM} \frac{dt}{t} \right\}^{1/2} \\ &\sim 2^{-2ikM} \left\{ \iint_{R_Q} |A(x, t)|^2 \frac{dx dt}{t} \right\}^{1/2} \lesssim 2^{-i[2kM - n(1/p - 1/2)]} |2^i Q|^{1/p - 1/2} \sim 2^{-i\gamma_2} |2^i Q|^{1/p - 1/2}, \end{aligned}$$

where $\gamma_2 := 2kM - n(1/p - 1/2) > 0$.

For I_2 , via some similar calculations to the estimate of I_1 , we see that

$$\begin{aligned} I_2 &\lesssim \int_0^{l(Q)} \left[\int_t^{2^i l(Q)} \left(\frac{t}{s} \right)^{4kb} \left\{ \frac{s^{2k}}{[2^i l(Q)]^{2k}} \right\}^{2M} \frac{ds}{s} \right]^{1/2} \|A(\cdot, t)\|_{L^2(\mathbb{R}^n)} \frac{dt}{t} \\ &\lesssim \int_0^{l(Q)} \left\{ \left[\frac{t}{[2^i l(Q)]} \right]^{2kb} + \left[\frac{t}{[2^i l(Q)]} \right]^{2kM} \right\} \|A(\cdot, t)\|_{L^2(\mathbb{R}^n)} \frac{dt}{t} \\ &\lesssim \left\{ \iint_{R_Q} |A(x, t)|^2 \frac{dx dt}{t} \right\}^{1/2} \left\{ \int_0^{l(Q)} \left(\left[\frac{t}{[2^i l(Q)]} \right]^{4kb} + \left[\frac{t}{[2^i l(Q)]} \right]^{4kM} \right) \frac{dt}{t} \right\}^{1/2} \\ &\lesssim (2^{-2ikb} + 2^{-2ikM}) |Q|^{1/2 - 1/p} \sim (2^{-i\gamma_1} + 2^{-i\gamma_2}) |2^i Q|^{1/2 - 1/p}. \end{aligned}$$

Combining the estimates of I_1 and I_2 , we obtain

$$O \lesssim (2^{-i\gamma_1} + 2^{-i\gamma_2}) |2^i Q|^{1/2 - 1/p}. \quad (5.13)$$

By (5.12) and (5.13), we conclude that

$$\|Q^f A\|_{T^p(\mathbb{R}_+^{n+1})}^p \lesssim \|\chi_{2R_Q} Q^f A\|_{T^p(\mathbb{R}_+^{n+1})}^p + \sum_{i=2}^{\infty} \|\chi_{S_i(R_Q)} Q^f A\|_{T^p(\mathbb{R}_+^{n+1})}^p \lesssim 1 + \sum_{i=2}^{\infty} (2^{-i\gamma_1 p} + 2^{-i\gamma_2 p}) \lesssim 1.$$

Thus, (5.10) holds, which completes the proof of Lemma 5.5. \square

As an application of Lemma 5.5, we obtain the following boundedness of $Q_{\psi, L}$ and $\pi_{\psi, L}$.

Lemma 5.6. *Let $p \in (0, 1]$, $\omega \in [0, \pi/2)$, L be the operator of type ω satisfying the assumptions (A_1) , (A_2) and (A_3) in Section 2, $\alpha \in (0, \infty)$, $\beta \in (n(1/p - 1/2)/(2k), \infty)$ and $\mu \in (\omega, \pi/2)$. Then*

- (i) *the operator $Q_{\psi, L}$, originally defined on $L^2(\mathbb{R}^n)$ as in (5.1) with $\psi \in \Psi_{\alpha, \beta}(S_\mu^0)$, can be extended to a bounded linear operator from $H_L^p(\mathbb{R}^n)$ to $T^p(\mathbb{R}_+^{n+1})$.*

- (ii) the operator $\pi_{\psi,L}$, originally defined on $T^2(\mathbb{R}_+^{n+1})$ as in (5.4) with $\psi \in \Psi_{\beta,\alpha}(S_\mu^0)$, can be extended to a bounded linear operator from $T^p(\mathbb{R}_+^{n+1})$ to $H_L^p(\mathbb{R}^n)$.

Proof. The proof of Lemma 5.6 is quite similar to that of [39, Proposition 4.9]. For the convenience of the reader, we present the details. We first recall a Calderón reproducing formula from [39, (4.12)]. For all $\psi \in \Psi(S_\mu^0)$, there exists a function $\tilde{\psi} \in \Psi(S_\mu^0)$ such that

$$\int_0^\infty \psi(t)\tilde{\psi}(t) \frac{dt}{t} = 1.$$

Moreover, we have

$$\pi_{\psi,L} \circ Q_{\tilde{\psi},L} = \pi_{\tilde{\psi},L} \circ Q_{\psi,L} = I \text{ in } L^2(\mathbb{R}^n). \quad (5.14)$$

In particular, let $\psi_0(z) := ze^{-z}$ for all $z \in S_\mu^0$. We then choose $\tilde{\psi}_0(z) := C(M)z^M e^{-z}$ for all $z \in S_\mu^0$ such that $\tilde{\psi}_0 \in \Psi_{M,N}(S_\mu^0)$ for any $N \in (0, \infty)$, where M is the smallest positive integer larger than $n(1/p - 1/2)/(2k)$ and $C(M) \int_0^\infty t^M e^{-2t} dt = 1$.

By Definition 5.1 and (5.1), we see that for all $f \in H_L^p(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$,

$$\|Q_{\psi_0,L}f\|_{T^p(\mathbb{R}_+^{n+1})} = \|f\|_{H_L^p(\mathbb{R}^n)},$$

which implies that $Q_{\psi_0,L}$ is bounded from $H_L^p(\mathbb{R}^n)$ to $T^p(\mathbb{R}_+^{n+1})$. For all $\psi \in \Psi_{\alpha,\beta}(S_\mu^0)$, by this, together with the Calderón reproducing formula (5.14) and Lemma 5.5 with $f := 1$ therein, we conclude that for all $f \in H_L^p(\mathbb{R}^n)$,

$$\|Q_{\psi,L}f\|_{T^p(\mathbb{R}_+^{n+1})} \sim \|Q_{\psi,L} \circ \pi_{\tilde{\psi}_0,L} \circ Q_{\psi_0,L}f\|_{T^p(\mathbb{R}_+^{n+1})} \lesssim \|Q_{\psi_0,L}f\|_{T^p(\mathbb{R}_+^{n+1})} \sim \|f\|_{H_L^p(\mathbb{R}^n)}.$$

That is, $Q_{\psi,L}$ is bounded from $H_L^p(\mathbb{R}^n)$ to $T^p(\mathbb{R}_+^{n+1})$, which completes the proof of (i).

On the other hand, for all $\psi \in \Psi_{\beta,\alpha}(S_\mu^0)$, since $\psi_0 \in \Psi_{1,\beta}(S_\mu^0)$, it follows from Lemma 5.5 with $f := 1$ therein that for all $F \in T^p(\mathbb{R}_+^{n+1}) \cap T^2(\mathbb{R}_+^{n+1})$,

$$\|\pi_{\psi,L}F\|_{H_L^p(\mathbb{R}^n)} = \|Q_{\psi_0,L} \circ \pi_{\psi,L}F\|_{T^p(\mathbb{R}_+^{n+1})} \lesssim \|F\|_{T^p(\mathbb{R}_+^{n+1})},$$

which shows that $\pi_{\psi,L}$ is bounded from $T^p(\mathbb{R}_+^{n+1})$ to $H_L^p(\mathbb{R}^n)$. This finishes the proof of (ii) and hence Lemma 5.6. \square

Proof of Theorem 5.2. By Definitions 4.1 and 5.1, to show Theorem 5.2, it suffices to prove that $\mathbb{H}_L^p(\mathbb{R}^n) = \mathbb{H}_{\psi,L}^p(\mathbb{R}^n)$ with equivalent norms.

The inclusion $\mathbb{H}_L^p(\mathbb{R}^n) \subset \mathbb{H}_{\psi,L}^p(\mathbb{R}^n)$ is an easy consequence of the boundedness of $Q_{\psi,L}$ from $H_L^p(\mathbb{R}^n)$ to $T^p(\mathbb{R}_+^{n+1})$, which is true by Lemma 5.6(i). We now prove $\mathbb{H}_{\psi,L}^p(\mathbb{R}^n) \subset \mathbb{H}_L^p(\mathbb{R}^n)$. Let $\psi_0(z) := ze^z$ for all $z \in S_\mu^0$. Observe that for any $\psi \in \Psi_{\alpha,\beta}(S_\mu^0)$, we can choose $\tilde{\psi}(z) := \tilde{C}(M)z^M e^{-z}$ for all $z \in S_\mu^0$ such that (5.14) holds, where $\tilde{C}(M)$ is a constant such that $\tilde{C}(M) \int_0^\infty t^{M-1} e^{-t} \psi(t) dt = 1$. From (5.14), Lemma 5.5 with $f := 1$ therein, and Lemma 5.6(i), we infer that for all $f \in \mathbb{H}_{\psi,L}^p(\mathbb{R}^n)$,

$$\|f\|_{H_L^p(\mathbb{R}^n)} = \|Q_{\psi_0,L}f\|_{T^p(\mathbb{R}_+^{n+1})} = \|Q_{\psi_0,L} \circ \pi_{\tilde{\psi},L} \circ Q_{\psi,L}f\|_{T^p(\mathbb{R}_+^{n+1})} \lesssim \|Q_{\psi,L}f\|_{T^p(\mathbb{R}_+^{n+1})} \sim \|f\|_{\mathbb{H}_{\psi,L}^p(\mathbb{R}^n)},$$

which implies that $\mathbb{H}_{\psi,L}^p(\mathbb{R}^n) \subset \mathbb{H}_L^p(\mathbb{R}^n)$. This finishes the proof of Theorem 5.2. \square

6 Riesz transforms on $H_{L_i}^p(\mathbb{R}^n)$ for $i \in \{1, 2\}$

In this section, for the $2k$ -order divergence form homogeneous elliptic operator L_1 with complex bounded measurable coefficients and the $2k$ -order Schrödinger type operator L_2 , we consider the behavior of their Riesz transforms $\nabla^k L_i^{-1/2}$ on the Hardy space $H_{L_i}^p(\mathbb{R}^n)$, respectively for $i \in \{1, 2\}$. First, we study the boundedness of $\nabla^k L_i^{-1/2}$ on $H_{L_i}^p(\mathbb{R}^n)$ for $i \in \{1, 2\}$. To this end, we need the following useful estimates.

Lemma 6.1. Let $p \in (0, 1]$, $M, k \in \mathbb{N}$, L_1 be the $2k$ -order divergence form homogenous elliptic operator with complex bounded measurable coefficients and L_2 the $2k$ -order Schrödinger type operator. Then, there exists a positive constant C such that for all $i \in \{1, 2\}$, closed sets E, F in \mathbb{R}^n with $\text{dist}(E, F) > 0$, $f \in L^2(\mathbb{R}^n)$ supported in E and $t \in (0, \infty)$,

$$\left\| \nabla^k L_i^{-1/2} (I - e^{-tL_i})^M f \right\|_{L^2(F)} \leq C \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^M \|f\|_{L^2(E)} \quad (6.1)$$

and

$$\left\| \nabla^k L_i^{-1/2} (tL_i e^{-tL_i})^M f \right\|_{L^2(F)} \leq C \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^M \|f\|_{L^2(E)}. \quad (6.2)$$

Proof. We prove this lemma by borrowing some ideas from [36]. Let $i \in \{1, 2\}$. From [6, Theorem 1.1] and [47, Theorem 8.1], we deduce that $\nabla^k L_i^{-1/2}$ is bounded on $L^2(\mathbb{R}^n)$. Thus, it suffices to prove Lemma 6.1 in the case that $t < [\text{dist}(E, F)]^{2k}$. By the H_∞ functional calculus in $L^2(\mathbb{R}^n)$, we know that for all $f \in L^2(\mathbb{R}^n)$,

$$L_i^{-1/2} f = \frac{1}{2\sqrt{\pi}} \int_0^\infty e^{-sL_i} s^{-1/2} f \, ds, \quad (6.3)$$

which, together with the change of variables, yields that

$$\begin{aligned} \nabla^k L_i^{-1/2} (I - e^{-tL_i})^M f &= \frac{1}{2\sqrt{\pi}} \int_0^\infty \nabla^k e^{-sL_i} (I - e^{-tL_i})^M f \frac{ds}{\sqrt{s}} \\ &= \frac{\sqrt{M+2}}{2\sqrt{\pi}} \int_0^\infty \nabla^k e^{-(M+2)sL_i} (I - e^{-tL_i})^M f \frac{ds}{\sqrt{s}} \\ &= \frac{\sqrt{M+2}}{2\sqrt{\pi}} \int_0^t \sqrt{s} \nabla^k e^{-(M+2)sL_i} \left[\sum_{j=0}^M \binom{M}{j} (-1)^j e^{-jtL_i} \right] f \frac{ds}{s} \\ &\quad + \frac{\sqrt{M+2}}{2\sqrt{\pi}} \int_t^\infty \sqrt{s} \nabla^k e^{-(M+2)sL_i} (I - e^{-tL_i})^M f \frac{ds}{s} \\ &=: \text{I} + \text{O}, \end{aligned}$$

where $\binom{M}{j}$ denotes the binomial coefficient.

To estimate I, we write

$$\begin{aligned} \text{I} &= \frac{\sqrt{M+2}}{2\sqrt{\pi}} \int_0^t \sqrt{s} \nabla^k e^{-sL_i} e^{-(M+1)sL_i} f \frac{ds}{s} + \sum_{j=1}^M \frac{\sqrt{M+2}}{2\sqrt{\pi}} \binom{M}{j} (-1)^j \int_0^t \nabla^k e^{-jtL_i} e^{-(M+2)sL_i} f \frac{ds}{\sqrt{s}} \\ &=: \text{I}_0 + \sum_{j=1}^M \text{I}_j. \end{aligned}$$

For I_0 , it follows, from Minkowski's inequality, Propositions 3.4 and 3.6, Lemma 3.2 and the assumption $t < [\text{dist}(E, F)]^{2k}$, that

$$\begin{aligned} \|\text{I}_0\|_{L^2(F)} &\lesssim \int_0^t \left\| \sqrt{s} \nabla^k e^{-sL_i} e^{-(M+1)sL_i} f \right\|_{L^2(F)} \frac{ds}{s} \\ &\lesssim \int_0^t \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}} \right\} \|f\|_{L^2(E)} \frac{ds}{s} \\ &\lesssim \exp \left\{ -\frac{\tilde{C}_1 [\text{dist}(E, F)]^{2k/(2k-1)}}{t^{1/(2k-1)}} \right\} \int_0^t \exp \left\{ -\frac{\tilde{C}_2 [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}} \right\} \frac{ds}{s} \|f\|_{L^2(E)} \end{aligned}$$

$$\begin{aligned} &\lesssim \exp \left\{ -\frac{\tilde{C}_1 [\text{dist}(E, F)]^{2k/(2k-1)}}{t^{1/(2k-1)}} \right\} \frac{t}{[\text{dist}(E, F)]^{2k}} \|f\|_{L^2(E)} \\ &\lesssim \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^M \|f\|_{L^2(E)}, \end{aligned}$$

where \tilde{C} , \tilde{C}_1 , \tilde{C}_2 are positive constants such that $\tilde{C}_1 + \tilde{C}_2 = \tilde{C}$.

For each I_j , $j \geq 1$, by Lemma 3.2 and Propositions 3.4 and 3.6, we see that

$$\begin{aligned} \|I_j\|_{L^2(F)} &\lesssim \frac{1}{\sqrt{t}} \int_0^t \left\| \left(\sqrt{jt} \nabla^k e^{-jtL_i} \right) \circ \left(e^{-(M+2)sL_i} \right) f \right\|_{L^2(F)} \frac{ds}{\sqrt{s}} \\ &\lesssim \frac{1}{\sqrt{t}} \|f\|_{L^2(E)} \int_0^t \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{t^{1/(2k-1)}} \right\} \frac{ds}{\sqrt{s}} \\ &\lesssim \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^M \|f\|_{L^2(E)}, \end{aligned}$$

which, together with the estimate of I_0 , implies that

$$\|I\|_{L^2(F)} \lesssim \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^M \|f\|_{L^2(E)}, \quad (6.4)$$

here and in what follows, \tilde{C} always denotes a positive constant. We now estimate O by writing

$$\begin{aligned} O &\sim \int_t^\infty \sqrt{s} \nabla^k e^{-sL_i} e^{-MsL_i} (I - e^{-tL_i})^M e^{-sL_i} f \frac{ds}{s} \\ &\sim \int_t^\infty (\sqrt{s} \nabla^k e^{-sL_i}) \circ (e^{-sL_i} - e^{-(s+t)L_i})^M \circ (e^{-sL_i}) f \frac{ds}{s}. \end{aligned}$$

Using the analytic property of semigroups and Lemma 3.1, we conclude that for all $g \in L^2(\mathbb{R}^n)$ supported in the closed set E and $t < s$,

$$\begin{aligned} \left\| [e^{-sL_i} - e^{-(s+t)L_i}] g \right\|_{L^2(F)} &= \left\| - \int_0^t \frac{\partial}{\partial r} (e^{-(s+r)L_i}) g dr \right\|_{L^2(F)} \\ &\lesssim \int_0^t \left\| (s+r)L_i e^{-(s+r)L_i} g \right\|_{L^2(F)} \frac{dr}{s+r} \\ &\lesssim \int_0^t \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}} \right\} \frac{dr}{s+r} \|g\|_{L^2(E)} \\ &\lesssim \frac{t}{s} \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}} \right\} \|g\|_{L^2(E)}. \end{aligned}$$

Thus,

$$\left\| \frac{s}{t} [e^{-sL_i} - e^{-(s+t)L_i}] g \right\|_{L^2(F)} \lesssim \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}} \right\} \|g\|_{L^2(E)}. \quad (6.5)$$

Therefore, from Minkowski's inequality, (6.5), Lemma 3.2, Propositions 3.4 and 3.6, and the change of variables, we deduce that

$$\|O\|_{L^2(F)} \lesssim \int_t^\infty \left\| (\sqrt{s} \nabla^k e^{-sL_i}) \circ \left(\frac{s}{t} [e^{-sL_i} - e^{-(s+t)L_i}] \right)^M \circ (e^{-sL_i}) f \right\|_{L^2(F)} \left(\frac{t}{s} \right)^M \frac{ds}{s}$$

$$\begin{aligned} &\lesssim \|f\|_{L^2(E)} \int_t^\infty \left(\frac{t}{s}\right)^M \exp\left\{-\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}}\right\} \frac{ds}{s} \\ &\lesssim \left(\frac{[\text{dist}(E, F)]^{2k}}{t}\right)^{-M} \|f\|_{L^2(E)}. \end{aligned}$$

Combining this estimate with (6.4), we have

$$\left\| \nabla^k L_i^{-1/2} (I - e^{-tL_i})^M f \right\|_{L^2(F)} \lesssim \left(\frac{[\text{dist}(E, F)]^{2k}}{t}\right)^{-M} \|f\|_{L^2(E)},$$

that is, (6.1) holds.

Now, we prove (6.2). Using (6.3) and the change of variables, we see that

$$\begin{aligned} \nabla^k L_i^{-1/2} (tL_i e^{-tL_i})^M f &= \frac{1}{2\sqrt{\pi}} \int_0^\infty \nabla^k e^{-sL_i} (tL_i e^{-tL_i})^M f \frac{ds}{\sqrt{s}} \\ &\sim \int_0^\infty \nabla^k e^{-(M+1)sL_i} (tL_i e^{-tL_i})^M f \frac{ds}{\sqrt{s}} \\ &\sim \int_0^t \nabla^k e^{-(M+1)sL_i} (tL_i e^{-tL_i})^M f \frac{ds}{\sqrt{s}} + \int_t^\infty \dots \\ &=: B + D. \end{aligned}$$

By an application of the analytic property of semigroups, Propositions 3.4 and 3.6, and Lemmas 3.1 and 3.2, we conclude that

$$\begin{aligned} \|B\|_{L^2(F)} &\lesssim \frac{1}{\sqrt{t}} \int_0^t \left\| \left(\sqrt{\frac{t}{2}} \nabla^k e^{-\frac{1}{2}L_i} \right) \circ \left(e^{-(M+1)sL_i} \right) \circ \left(\frac{t}{2} L_i e^{-\frac{1}{2}L_i} \right) \circ (tL_i e^{-tL_i})^{M-1} f \right\|_{L^2(F)} \frac{ds}{\sqrt{s}} \\ &\lesssim \frac{1}{\sqrt{t}} \int_0^t \exp\left\{-\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{t^{1/(2k-1)}}\right\} \|f\|_{L^2(E)} \frac{ds}{\sqrt{s}} \\ &\lesssim \exp\left\{-\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{t^{1/(2k-1)}}\right\} \|f\|_{L^2(E)} \lesssim \left(\frac{t}{[\text{dist}(E, F)]^{2k}}\right)^M \|f\|_{L^2(E)}, \end{aligned}$$

where C is a positive constant.

For the estimate of D , similar to the estimate for B , we write

$$D = \int_t^\infty (\sqrt{s} \nabla^k e^{-sL_i}) \circ \left(\frac{t}{s}\right)^M \circ [sL_i e^{-(s+t)L_i}]^M f \frac{ds}{s}$$

and we estimate $sL_i e^{-(s+t)L_i} f$ by

$$\begin{aligned} \|sL_i e^{-(s+t)L_i} f\|_{L^2(F)} &= \left\| \frac{s}{t} e^{-sL_i} \int_0^t \frac{\partial}{\partial r} (rL_i e^{-rL_i}) f dr \right\|_{L^2(F)} \\ &\lesssim \left\| \frac{s}{t} e^{-sL_i} \int_0^t [L_i e^{-rL_i} f - rL_i^2 e^{-rL_i} f] dr \right\|_{L^2(F)} \\ &\lesssim \left\| \frac{s}{t} e^{-sL_i} \int_0^t L_i e^{-rL_i} f dr \right\|_{L^2(F)} + \left\| \frac{s}{t} e^{-sL_i} \int_0^t rL_i^2 e^{-rL_i} f dr \right\|_{L^2(F)} \\ &=: V_1 + V_2. \end{aligned}$$

By Minkowski's inequality, Lemma 3.1 and $r < t < s$, we conclude that

$$V_1 \lesssim \frac{s}{t} \int_0^t \|L_i e^{-(s+r)L_i} f\|_{L^2(F)} dr \lesssim \frac{s}{t} \int_0^t \exp\left\{-\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{(s+r)^{1/(2k-1)}}\right\} \|f\|_{L^2(E)} \frac{dr}{s}$$

$$\lesssim \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}} \right\} \|f\|_{L^2(E)}.$$

Similarly, we see that

$$\begin{aligned} V_2 &\lesssim \frac{s}{t} \int_0^t \left\| [(r+s)L_i]^2 e^{-(r+s)L_i} f \right\|_{L^2(F)} \frac{dr}{r+s} \\ &\lesssim \frac{s}{t} \|f\|_{L^2(E)} \int_0^t \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{(r+s)^{1/(2k-1)}} \right\} \frac{dr}{r+s} \\ &\lesssim \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}} \right\} \|f\|_{L^2(E)}, \end{aligned}$$

which, together with the estimate of V_1 , shows that the family $\{sL_i e^{-(s+t)L_i}\}_{t>0}$ of operators satisfies the k -Davies-Gaffney estimate in s . Thus, from Minkowski's inequality, Lemmas 3.1 and 3.2, Propositions 3.4 and 3.6, and the change of variables, we deduce that

$$\begin{aligned} \|D\|_{L^2(F)} &\lesssim \int_t^\infty \left\| \left(\sqrt{s} \nabla^k e^{-sL_i} \right) \circ \left(\frac{t}{s} \right)^M \circ \left(sL_i e^{-(s+t)L_i} \right)^M f \right\|_{L^2(F)} \frac{ds}{s} \\ &\lesssim \|f\|_{L^2(E)} \int_t^\infty \left(\frac{t}{s} \right)^M \exp \left\{ -\frac{\tilde{C} [\text{dist}(E, F)]^{2k/(2k-1)}}{s^{1/(2k-1)}} \right\} \frac{ds}{s} \\ &\lesssim \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^M \|f\|_{L^2(E)}. \end{aligned}$$

Combining the estimates for B and D , we see that

$$\left\| \nabla^k L_i^{-1/2} (tL_i e^{-tL_i})^M f \right\|_{L^2(F)} \lesssim \left(\frac{t}{[\text{dist}(E, F)]^{2k}} \right)^M \|f\|_{L^2(E)},$$

which shows that (6.2) also holds. This finishes the proof of Lemma 6.1. \square

With the help of Lemma 6.1, we show that the Riesz transform $\nabla^k(L_i^{-1/2})$ is bounded from $H_L^p(\mathbb{R}^n)$ to the classical Hardy space $H^p(\mathbb{R}^n)$, which when $p = 1$, $i = 2$ and $k = 1$ was first obtained in [35].

Theorem 6.2. *Let $k \in \mathbb{N}$, $p \in (n/(n+k), 1]$, L_1 be the $2k$ -order divergence form homogeneous elliptic operator with complex bounded measurable coefficients and L_2 the $2k$ -order Schrödinger type operator. Then, for all $i \in \{1, 2\}$, the Riesz transform $\nabla^k(L_i^{-1/2})$ is bounded from $H_{L_i}^p(\mathbb{R}^n)$ to the classical Hardy space $H^p(\mathbb{R}^n)$.*

Proof. Let $i \in \{1, 2\}$. We first claim that to prove Theorem 6.2, it suffices to show that $\nabla^k(L_i^{-1/2})$ maps each $(H_{L_i}^p, \epsilon, M)$ -molecule m as in Definition 4.3 with $\epsilon > 0$ and $M > n(1/p - 1/2)/(2k)$ into a classical $H^p(\mathbb{R}^n)$ -molecule in [50] up to a harmless constant multiple.

Indeed, assume this claim for the moment. For any $f \in \mathbb{H}_{L_i}^p(\mathbb{R}^n)$, by Theorem 4.5, there exist $\{\lambda_j\}_{j=0}^\infty \in l^p$ and a sequence $\{m_j\}_{j=0}^\infty$ of $(H_{L_i}^p, \epsilon, M)$ -molecules such that $f = \sum_{j=0}^\infty \lambda_j m_j$ is a molecular $(H_L^p, 2, \epsilon, M)$ -representation of f and

$$\|f\|_{H_{L_i}^p(\mathbb{R}^n)} \sim \left(\sum_{j=0}^\infty |\lambda_j|^p \right)^{1/p}.$$

Moreover, from the $L^2(\mathbb{R}^n)$ -boundedness of $\nabla^k(L_i^{-1/2})$ and the fact that $f = \sum_{j=0}^\infty \lambda_j m_j$ holds in $L^2(\mathbb{R}^n)$, it follows that

$$\nabla^k(L_i^{-1/2})f = \nabla^k(L_i^{-1/2}) \left(\sum_{j=0}^\infty \lambda_j m_j \right) = \sum_{j=0}^\infty \lambda_j \nabla^k(L_i^{-1/2})m_j \quad (6.6)$$

in $L^2(\mathbb{R}^n)$ and hence in the space $\mathcal{S}'(\mathbb{R}^n)$ of Schwartz distributions, which, together with the above claim, implies that (6.6) is a classical molecular decomposition of $\nabla^k(L_i^{-1/2})f$ in $H^p(\mathbb{R}^n)$. Thus, by the molecular characterization of $H^p(\mathbb{R}^n)$ in [50], we further conclude that

$$\left\| \nabla^k(L_i^{-1/2})f \right\|_{H^p(\mathbb{R}^n)} \lesssim \left(\sum_{j=0}^{\infty} |\lambda_j|^p \right)^{1/p} \sim \|f\|_{H_{L_i}^p(\mathbb{R}^n)},$$

which, combined with a density argument, then shows that $\nabla^k(L_i^{-1/2})$ is bounded from $H_{L_i}^p(\mathbb{R}^n)$ to $H^p(\mathbb{R}^n)$.

Let m be an $(H_{L_i}^p, \epsilon, M)$ -molecule associated with the cube Q as in Definition 4.3 with $\epsilon \in (0, \infty)$ and $M > n(1/p - 1/2)/(2k)$. To prove the above claim, we need to prove that $\nabla^k(L_i^{-1/2})m$ is a classical $H^p(\mathbb{R}^n)$ -molecule in [50] up to a harmless constant multiple. To this end, we only need to show that $\nabla^k(L_i^{-1/2})m$ is a following defined $H^p(\mathbb{R}^n)$ -molecule in [39, 40], from which it follows that it is also a classical molecule in [50]. In what follows, for any $\gamma \in \mathbb{R}$, we denote by $\lfloor \gamma \rfloor$ the maximal integer not more than γ . Let $p \in (0, 1]$ and Q be a cube in \mathbb{R}^n . A function $\tilde{m} \in L^2(\mathbb{R}^n)$ is called an $H^p(\mathbb{R}^n)$ -molecule associated with Q if there exists a positive constant $\epsilon \in (0, \infty)$ such that

(i) for all $j \in \mathbb{Z}_+$,

$$\|\tilde{m}\|_{L^2(S_j(Q))} \lesssim [2^j l(Q)]^{n(1/p-1/2)} 2^{-j\epsilon}; \quad (6.7)$$

(ii) there exists a non-negative integer $M \in \mathbb{Z}_+$ with $M \geq \lfloor n(1/p - 1) \rfloor$ such that for all multi-indices α with $0 \leq |\alpha| \leq M$,

$$\int_{\mathbb{R}^n} x^\alpha \tilde{m}(x) dx = 0. \quad (6.8)$$

We first prove that $\nabla^k(L_i^{-1/2})m$ satisfies (6.7). For all $j \in \{0, 1, 2\}$, by the $L^2(\mathbb{R}^n)$ -boundedness of $\nabla^k(L_i^{-1/2})$ and (4.5), we see that

$$\left\| \nabla^k(L_i^{-1/2})m \right\|_{L^2(S_j(Q))} \lesssim \left\| \nabla^k(L_i^{-1/2})m \right\|_{L^2(\mathbb{R}^n)} \lesssim \|m\|_{L^2(\mathbb{R}^n)} \lesssim |Q|^{1/2-1/p}.$$

When $j > 2$, we write

$$\begin{aligned} \left\| \nabla^k(L_i^{-1/2})m \right\|_{L^2(S_j(Q))} &\leq \left\| \nabla^k(L_i^{-1/2}) \left(I - e^{-[l(Q)]^{2k} L_i} \right)^M m \right\|_{L^2(S_j(Q))} \\ &\quad + \left\| \nabla^k(L_i^{-1/2}) \left[I - \left(I - e^{-[l(Q)]^{2k} L_i} \right)^M \right] m \right\|_{L^2(S_j(Q))} \\ &=: \text{I} + \text{O}. \end{aligned}$$

From an application of Lemma 6.1 and (4.5), it follows that

$$\begin{aligned} \text{I} &\lesssim \left\| \nabla^k(L_i^{-1/2}) \left(I - e^{-[l(Q)]^{2k} L_i} \right)^M (m \chi_{2^{j-2}Q}) \right\|_{L^2(S_j(Q))} \\ &\quad + \left\| \nabla^k(L_i^{-1/2}) \left(I - e^{-[l(Q)]^{2k} L_i} \right)^M (m \chi_{\mathbb{R}^n \setminus (2^{j+1}Q)}) \right\|_{L^2(S_j(Q))} \\ &\quad + \left\| \nabla^k(L_i^{-1/2}) \left(I - e^{-[l(Q)]^{2k} L_i} \right)^M (m \chi_{2^{j+1}Q \setminus (2^{j-2}Q)}) \right\|_{L^2(S_j(Q))} \\ &\lesssim \left[\frac{\text{dist}(S_j(Q), 2^{j-2}Q)}{l(Q)} \right]^{2kM} \|m \chi_{2^{j-2}Q}\|_{L^2(\mathbb{R}^n)} \end{aligned}$$

$$\begin{aligned}
& + \left[\frac{\text{dist}(S_j(Q), \mathbb{R}^n \setminus (2^{j+1}Q))}{l(Q)} \right]^{2kM} \|m\chi_{\mathbb{R}^n \setminus (2^{j+1}Q)}\|_{L^2(\mathbb{R}^n)} + \|m\chi_{2^{j+1}Q \setminus (2^{j-1}Q)}\|_{L^2(\mathbb{R}^n)} \\
& \lesssim 2^{-2jkM} [l(Q)]^{n(1/2-1/p)} + [2^j l(Q)]^{n(1/2-1/p)} 2^{-j\epsilon}.
\end{aligned}$$

Let $\tilde{\epsilon} := \min\{\epsilon, 2kM - n(1/p - 1/2)\} > 0$. We then have

$$I \lesssim [2^j l(Q)]^{n(1/2-1/p)} 2^{-j\tilde{\epsilon}}. \quad (6.9)$$

To estimate O , from Lemma 6.1 and (4.2), we deduce that

$$\begin{aligned}
O & \lesssim \sup_{1 \leq \ell \leq M} \left\| \nabla^k L_i^{-1/2} e^{-\ell[l(Q)]^{2k} L_i} m \right\|_{L^2(S_j(Q))} \\
& \sim \sup_{1 \leq \ell \leq M} \left\| \nabla^k L_i^{-1/2} \left(\frac{\ell}{M} [l(Q)]^{2k} L_i e^{-\frac{\ell}{M} [l(Q)]^{2k} L_i} \right)^M ([l(Q)]^{-2k} L_i^{-1})^M m \right\|_{L^2(S_j(Q))} \\
& \sim \sup_{1 \leq \ell \leq M} \left\| \nabla^k L_i^{-1/2} \left(\frac{\ell}{M} [l(Q)]^{2k} L_i e^{-\frac{\ell}{M} [l(Q)]^{2k} L_i} \right)^M \left[\chi_{2^{j-2}Q} ([l(Q)]^{-2k} L_i^{-1})^M \right] m \right\|_{L^2(S_j(Q))} \\
& \quad + \sup_{1 \leq \ell \leq M} \left\| \nabla^k L_i^{-1/2} \left(\frac{\ell}{M} [l(Q)]^{2k} L_i e^{-\frac{\ell}{M} [l(Q)]^{2k} L_i} \right)^M \left[\chi_{\mathbb{R}^n \setminus 2^{j+1}Q} ([l(Q)]^{-2k} L_i^{-1})^M \right] m \right\|_{L^2(S_j(Q))} \\
& \quad + \sup_{1 \leq \ell \leq M} \left\| \nabla^k L_i^{-1/2} \left(\frac{\ell}{M} [l(Q)]^{2k} L_i e^{-\frac{\ell}{M} [l(Q)]^{2k} L_i} \right)^M \left[\chi_{2^{j+1}Q \setminus 2^{j-2}Q} ([l(Q)]^{-2k} L_i^{-1})^M \right] m \right\|_{L^2(S_j(Q))} \\
& \lesssim 2^{-2jkM} \left\| ([l(Q)]^{-2k} L_i^{-1})^M m \right\|_{L^2(\mathbb{R}^n)} + \left\| \chi_{2^{j+1}Q \setminus 2^{j-2}Q} ([l(Q)]^{-2k} L_i^{-1})^M m \right\|_{L^2(\mathbb{R}^n)} \\
& \lesssim 2^{-2jkM} \left\{ \sum_{\tilde{k}=0}^{\infty} \left\| ([l(Q)]^{-2k} L_i^{-1})^M m \right\|_{L^2(S_{\tilde{k}}(Q))}^2 \right\}^{1/2} + [2^j l(Q)]^{n(1/2-1/p)} 2^{-j\epsilon} \\
& \lesssim 2^{-2jkM} \left\{ \sum_{\tilde{k}=0}^{\infty} 2^{-2\tilde{k}[\epsilon+n(1/p-1/2)]} \right\}^{1/2} [l(Q)]^{n(1/2-1/p)} + [2^j l(Q)]^{n(1/2-1/p)} 2^{-j\epsilon} \\
& \lesssim 2^{-j\tilde{\epsilon}} [2^j l(Q)]^{n/p-2/p},
\end{aligned}$$

which, together with (6.9), implies that $\nabla^k(L_i^{-1/2})m$ satisfies (6.7) with ϵ therein replaced by $\tilde{\epsilon}$.

Now, we prove that $\nabla^k(L_i^{-1/2})m$ satisfies (6.8) by borrowing some ideas from the proof of Theorem 7.4 in [43]. Let $D(\sqrt{L_i})$ be the domain of $\sqrt{L_i}$ and $R(L_i^{-1/2})$ the range of $L_i^{-1/2}$. From [6, 47], it follows that $D(\sqrt{L_i}) = D(\mathfrak{a}_i)$, where $D(\mathfrak{a}_i) \subset W^{k,2}(\mathbb{R}^n)$ is the domain of the sesquilinear form associated with L_i , which implies that $R(L_i^{-1/2}) \subset W^{k,2}(\mathbb{R}^n)$. Let $\{\varphi_j\}_{j=1}^{\infty} \subset C_c^{\infty}(\mathbb{R}^n)$ such that

- (i) $\sum_{j=1}^{\infty} \varphi_j(x) = 1$ for almost every $x \in \mathbb{R}^n$;
- (ii) for each $j \in \mathbb{N}$, there exists a ball $B_j \subset \mathbb{R}^n$ such that $\text{supp } \varphi_j \subset 2B_j$, $\varphi_j := 1$ on B_j and $0 \leq \varphi_j \leq 1$;
- (iii) there exists a positive constant C_{φ} such that for all $j \in \mathbb{N}$ and $x \in \mathbb{R}^n$,

$$\sum_{\ell=1}^k |\nabla^{\ell} \varphi_j(x)| \leq C_{\varphi};$$

- (iv) there exists $N_{\varphi} \in \mathbb{N}$ such that $\sum_{j=1}^{\infty} \chi_{2B_j} \leq N_{\varphi}$.

For all $j \in \mathbb{N}$ and multi-indices α , let $\eta_j \in C_c^\infty(\mathbb{R}^n)$ such that $\eta_j := 1$ on $2B_j$ and $\text{supp } \eta_j \subset 4B_j$. Since $R(L_i^{-1/2}) \subset W^{k,2}(\mathbb{R}^n)$ and $\eta_j x^\alpha \in C_c^\infty(\mathbb{R}^n)$, we conclude that

$$\begin{aligned} \int_{\mathbb{R}^n} x^\alpha \nabla^k L_i^{-1/2} m(x) dx &= \int_{\mathbb{R}^n} x^\alpha \nabla^{k-1} \left(\sum_{j=1}^{\infty} \varphi_j \nabla L_i^{-1/2} \right) m(x) dx \\ &= \sum_{j=1}^{\infty} \int_{\mathbb{R}^n} x^\alpha \nabla^{k-1} \left(\varphi_j \nabla L_i^{-1/2} \right) m(x) dx \\ &= \sum_{j=1}^{\infty} \int_{\mathbb{R}^n} \eta_j x^\alpha \nabla^{k-1} \left(\nabla L_i^{-1/2} \right) m(x) dx \\ &= \sum_{j=1}^{\infty} (-1)^{k-1} \int_{\mathbb{R}^n} (\nabla^{k-1}(\eta_j x^\alpha)) \nabla(L_i^{-1/2}) m(x) dx. \end{aligned}$$

Thus, for all $|\alpha| \leq k-1 = n(1/[n/(n+k)]-1)$, we see that

$$\begin{aligned} \left| \int_{\mathbb{R}^n} x^\alpha \nabla^k L_i^{-1/2} m(x) dx \right| &\leq \sum_{j=1}^{\infty} \left| \int_{\mathbb{R}^n} (\nabla^{k-1}(\eta_j x^\alpha)) \nabla(L_i^{-1/2}) m(x) dx \right| \\ &\leq \sum_{j=1}^{\infty} \left| \int_{\mathbb{R}^n} \eta_j \nabla(L_i^{-1/2}) m(x) dx \right| \\ &= \sum_{j=1}^{\infty} \left| \int_{\mathbb{R}^n} \eta_j \nabla(\varphi_i L_i^{-1/2}) m(x) dx \right| \\ &= \sum_{j=1}^{\infty} \left| \int_{\mathbb{R}^n} \nabla(\eta_j) \varphi_i L_i^{-1/2} m(x) dx \right| = 0, \end{aligned}$$

which implies that $\nabla^k(L_i^{-1/2})m$ satisfies (6.8) with p and M respectively replaced by $n/(n+k)$ and $n(1/[n/(n+k)]-1)$. Thus, $\nabla^k(L_i^{-1/2})m$ is a classical $H^p(\mathbb{R}^n)$ molecule in [50], which completes the proof of Theorem 6.2. \square

On the Hardy space $H_{L_1}^p(\mathbb{R}^n)$, we further obtain its characterization by the Riesz transforms $\nabla^k(L_1^{-1/2})$. To this end, we first introduce some notions.

Definition 6.3. Let $p \in (0, 1]$ and L_1 be the $2k$ -order divergence form homogenous elliptic operator with complex bounded measurable coefficients. The *Riesz transform Hardy space* $H_{L_1, \text{Riesz}}^p(\mathbb{R}^n)$ is defined to be the completion of the set

$$\mathbb{H}_{L_1, \text{Riesz}}^p(\mathbb{R}^n) := \left\{ f \in L^2(\mathbb{R}^n) : \nabla^k(L_1^{-1/2})f \in H^p(\mathbb{R}^n) \right\}$$

with respect to the quasi-norm

$$\|f\|_{H_{L_1, \text{Riesz}}^p(\mathbb{R}^n)} := \left\| \nabla^k(L_1^{-1/2})f \right\|_{H^p(\mathbb{R}^n)}$$

for all $f \in \mathbb{H}_{L_1, \text{Riesz}}^p(\mathbb{R}^n)$.

We also need the following notion of $L^p - L^q$ k -off-diagonal estimates, which when $k = 1$ previously appeared in [3] (see also [39]).

Definition 6.4. Let $k \in \mathbb{N}$, $r, q \in (1, \infty)$ and $r \leq q$. A family $\{S_t\}_{t>0}$ of operators is said to satisfy the $L^r - L^q$ k -off-diagonal estimate, if there exist positive constant C and \tilde{C} such that for all closed sets $E, F \subset \mathbb{R}^n$ and $f \in L^r(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ supported in E ,

$$\|S_t f\|_{L^q(F)} \leq C t^{\frac{n}{2k}(\frac{1}{q} - \frac{1}{r})} \exp \left\{ -\tilde{C} \frac{[\text{dist}(E, F)]^{2k/(2k-1)}}{t^{1/(2k-1)}} \right\} \|f\|_{L^r(E)}.$$

On the $L^r - L^q$ k -off-diagonal estimate of the $2k$ -order divergence form homogeneous elliptic operator L_1 with complex bounded measurable coefficients, we have the following useful lemma.

Lemma 6.5. *Let L_1 be the $2k$ -order divergence form homogeneous elliptic operator with complex bounded measurable coefficients and $r \in (1, 2]$ such that the semigroup $\{e^{-tL_1}\}_{t>0}$ satisfies the $L^r - L^2$ k -off-diagonal estimate. Then the family $\{tL_1e^{-tL_1}\}_{t>0}$ of operators also satisfies the $L^r - L^2$ k -off-diagonal estimate.*

Proof. By the analytical property of $\{e^{-tL_1}\}_{t>0}$, we have $\{tL_1e^{-tL_1}\}_{t>0} = \{2(\frac{t}{2}L_1e^{-\frac{t}{2}L_1})(e^{-\frac{t}{2}L_1})\}_{t>0}$. Since the k -Davies-Gaffney estimate is just the $L^2 - L^2$ k -off-diagonal estimate, it follows, from Proposition 3.4 and Lemma 3.1, that $\{\frac{t}{2}L_1e^{-\frac{t}{2}L_1}\}_{t>0}$ satisfies the $L^2 - L^2$ k -off-diagonal estimate. Moreover, by the fact that $\{e^{-\frac{t}{2}L_1}\}_{t>0}$ satisfies the $L^r - L^2$ k -off-diagonal estimate and an argument similar to the proof of Lemma 3.2 with $\{A_t\}_{t>0}$ and $\{B_s\}_{s>0}$, respectively, replaced by $\{\frac{t}{2}L_1e^{-\frac{t}{2}L_1}\}_{t>0}$ and $\{e^{-\frac{t}{2}L_1}\}_{t>0}$, we conclude that $\{tL_1e^{-tL_1}\}_{t>0}$ also satisfies the $L^r - L^2$ k -off-diagonal estimate, which completes the proof of Lemma 6.5. \square

Proposition 6.6. *Let L_1 be the $2k$ -order divergence form homogeneous elliptic operator with complex bounded measurable coefficients and $r \in (1, 2]$ such that the semigroup $\{e^{-tL_1}\}_{t>0}$ satisfies the $L^r - L^2$ k -off-diagonal estimate. Then for all $p \in (0, 1]$ such that $p > rn/(n + kr)$ and $h \in \mathbb{H}_{L_1, \text{Riesz}}^p(\mathbb{R}^n)$,*

$$\|h\|_{H_{L_1}^p(\mathbb{R}^n)} \leq C \|\nabla^k L_1^{-1/2} h\|_{H^p(\mathbb{R}^n)}.$$

To prove Proposition 6.6, we need to recall some results concerning the homogenous Hardy-Sobolev space $\dot{H}^{k,p}(\mathbb{R}^n)$ (see, for example, [16, 34, 51, 52]).

Definition 6.7. Let $k \in \mathbb{N}$ and $p \in (0, 1]$. The *homogeneous Hardy-Sobolev space* $\dot{H}^{k,p}(\mathbb{R}^n)$ is defined to be the space

$$\dot{H}^{k,p}(\mathbb{R}^n) := \left\{ f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}_{k-1}(\mathbb{R}^n) : \|f\|_{\dot{H}^{k,p}(\mathbb{R}^n)} := \sum_{|\sigma|=k} \|\partial^\sigma f\|_{H^p(\mathbb{R}^n)} < \infty \right\},$$

where $\mathcal{S}'(\mathbb{R}^n)$ denotes the space of all Schwartz distributions on \mathbb{R}^n and $\mathcal{P}_{k-1}(\mathbb{R}^n)$ the class of all polynomials of order strictly less than k on \mathbb{R}^n .

Let $\ell \in \mathbb{N}$ be fixed. Let $\mathcal{S}(\mathbb{R}^n)$ denote the space of all Schwartz functions on \mathbb{R}^n and $\phi \in \mathcal{S}(\mathbb{R}^n)$ such that

- (i) ϕ is radial, $\text{supp } \phi \subset \{x \in \mathbb{R}^n : |x| < 1\}$ and for all $\xi \neq 0$, $\int_0^\infty |\widehat{\phi}(t\xi)|^2 \frac{dt}{t} = 1$, where $\widehat{\phi}$ denotes the Fourier transform of ϕ ,
- (ii) for all $|\gamma| \leq \ell$, $\int_{\mathbb{R}^n} x^\gamma \phi(x) dx = 0$.

For any given $\phi \in \mathcal{S}(\mathbb{R}^n)$ as above and all $f \in \mathcal{S}'(\mathbb{R}^n)$, let $Q_t f := \phi_t * f$, where $\phi_t := t^{-n} \phi(x/t)$ for all $t \in (0, \infty)$ and $x \in \mathbb{R}^n$. Let $p, q \in (0, \infty)$ and $\alpha \in \mathbb{R}$ such that $|\alpha| < \ell + 1$. The *homogeneous Triebel-Lizorkin space* $\dot{F}_{p,q}^\alpha(\mathbb{R}^n)$ is defined to be the space

$$\dot{F}_{p,q}^\alpha(\mathbb{R}^n) := \left\{ f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n) : \|f\|_{\dot{F}_{p,q}^\alpha(\mathbb{R}^n)} := \left\| \left\{ \int_0^\infty (t^{-\alpha} |Q_t f|)^q \frac{dt}{t} \right\}^{1/q} \right\|_{L^p(\mathbb{R}^n)} < \infty \right\},$$

where $\mathcal{P}(\mathbb{R}^n)$ denotes the class of all polynomials on \mathbb{R}^n (see, for example, [34, 51, 52]).

Let $\dot{W}^{k,2}(\mathbb{R}^n)$ for $k \in \mathbb{N}$ denote the *homogenous Sobolev space of order k* endowed with the norm $\|\cdot\|_{\dot{W}^{k,2}(\mathbb{R}^n)} := \|\nabla^k(\cdot)\|_{L^2(\mathbb{R}^n)}$. It is known that the homogeneous Sobolev space $\dot{W}^{k,2}(\mathbb{R}^n)$ and Hardy-Sobolev space $\dot{H}^{k,p}(\mathbb{R}^n)$ coincide, respectively, with the Triebel-Lizorkin space $\dot{F}_{2,2}^k(\mathbb{R}^n)$ and $\dot{F}_{p,2}^k(\mathbb{R}^n)$ with equivalent norms (see, for example, [51, p. 242]).

Definition 6.8. Let $k \in \mathbb{N}$, $\ell \geq k$ be any fixed positive integer and $p \in (0, 1]$. A function b is called an $\dot{H}^{k,p}(\mathbb{R}^n)$ -atom if it satisfies that

- (i) there exists a ball $B \subset \mathbb{R}^n$ such that $\text{supp } b \subset B$,
- (ii) for any $|\gamma| \leq \ell$, $\int_{\mathbb{R}^n} x^\gamma b(x) dx = 0$,
- (iii)

$$\|b\|_{\dot{F}_{2,2}^k(\mathbb{R}^n)} \leq |B|^{1/2-1/p}. \quad (6.10)$$

Lemma 6.9. Let $p \in (0, 1]$, $k \in \mathbb{N}$ and $f \in \dot{W}^{k,2}(\mathbb{R}^n) \cap \dot{H}^{k,p}(\mathbb{R}^n)$. Then there exist $\{\lambda_j\}_{j=0}^\infty \in l^p$ and a sequence $\{b_j\}_{j=0}^\infty$ of $\dot{H}^{k,p}(\mathbb{R}^n)$ -atoms such that $f = \sum_{j=0}^\infty \lambda_j b_j$ in $\dot{W}^{k,2}(\mathbb{R}^n) \cap \dot{H}^{k,p}(\mathbb{R}^n)$, and $\|f\|_{\dot{H}^{k,p}(\mathbb{R}^n)} \sim \{\sum_{j=0}^\infty |\lambda_j|^p\}^{1/p}$.

Proof. For any $f \in \dot{W}^{k,2}(\mathbb{R}^n) \cap \dot{H}^{k,p}(\mathbb{R}^n)$, by the coincidence of Sobolev spaces and Hardy-Sobolev spaces with Triebel-Lizorkin spaces, we know that $f \in \dot{F}_{2,2}^k(\mathbb{R}^n) \cap \dot{F}_{p,2}^k(\mathbb{R}^n)$. From this and a slight modification on the proof of [52, Proposition 4.3], together with the same observation as in Theorem 4.7 on the convergence of the atomic decomposition for elements in the tent spaces, we deduce all the desired conclusions of Lemma 6.9, which completes the proof of Lemma 6.9. \square

Proof of Proposition 6.6. For all $g \in L^2(\mathbb{R}^n)$, define the operator S_1 by setting, for all $x \in \mathbb{R}^n$,

$$S_1 g(x) := \left\{ \iint_{\Gamma(x)} \left| t^k \sqrt{L_1} e^{-t^{2k} L_1} g(y) \right|^2 \frac{dy dt}{t^{n+1}} \right\}^{1/2}.$$

For all $h \in \mathbb{H}_{\text{Riesz}, L_1}^p(\mathbb{R}^n)$, let $f := L_1^{-1/2} h$. Then $f \in \dot{W}^{k,2}(\mathbb{R}^n) \cap \dot{H}^{k,p}(\mathbb{R}^n)$ and, by Lemma 6.9, there exist $\{\lambda_j\}_{j=0}^\infty \in l^p$ and a sequence $\{b_j\}_{j=0}^\infty$ of $\dot{H}^{k,p}(\mathbb{R}^n)$ -atoms such that $f = \sum_{j=0}^\infty \lambda_j b_j$ in $\dot{W}^{k,2}(\mathbb{R}^n) \cap \dot{H}^{k,p}(\mathbb{R}^n)$ and, moreover, $(\sum_{j=0}^\infty |\lambda_j|^p)^{1/p} \sim \|f\|_{\dot{H}^{k,p}(\mathbb{R}^n)}$. By Theorem 5.2 with L replaced by L_1 , to show Proposition 6.6, we only need to prove that for all $f \in \dot{W}^{k,2}(\mathbb{R}^n) \cap \dot{H}^{k,p}(\mathbb{R}^n)$ with $p \in (nr/(n+kr), 1]$,

$$\left\| S_1 \sqrt{L_1} f \right\|_{L^p(\mathbb{R}^n)} \lesssim \|f\|_{\dot{H}^{k,p}(\mathbb{R}^n)}. \quad (6.11)$$

To prove (6.11), it suffices to prove that for all $\dot{H}^{k,p}(\mathbb{R}^n)$ -atoms b ,

$$\left\| S_1 \sqrt{L_1} b \right\|_{L^p(\mathbb{R}^n)} \lesssim 1. \quad (6.12)$$

Indeed, if (6.12) holds, by the $L^2(\mathbb{R}^n)$ -boundedness of S_1 which is deduced from (4.3), and [5, Theorem 1.1], we conclude that

$$\left\| S_1 \sqrt{L_1} f \right\|_{L^2(\mathbb{R}^n)} \lesssim \left\| \sqrt{L_1} f \right\|_{L^2(\mathbb{R}^n)} \sim \|\nabla^k f\|_{L^2(\mathbb{R}^n)} \sim \|f\|_{\dot{W}^{k,2}(\mathbb{R}^n)},$$

which, together with an argument similar to the proof of (5.11), yields that for almost every $x \in \mathbb{R}^n$, $|S_1 \sqrt{L_1} f(x)| \leq \sum_{j=0}^\infty |\lambda_j S_1 \sqrt{L_1} b_j(x)|$. This, combined with (6.12), shows that (6.11) is valid.

We now prove (6.12). For $j \in \mathbb{N}$, let $\mathcal{R}(S_j(Q)) := \cup_{x \in S_j(Q)} \Gamma(x)$ be the saw-tooth region based on $S_j(Q) \subset \mathbb{R}^n$. From Minkowski's inequality, Hölder's inequality and Fubini's theorem, we infer that

$$\begin{aligned} \left\| S_1 \sqrt{L_1} b \right\|_{L^p(\mathbb{R}^n)}^p &\lesssim \sum_{j=0}^\infty \left\| S_1 \sqrt{L_1} b \right\|_{L^p(S_j(Q))}^p \\ &\lesssim \left\| S_1 \sqrt{L_1} b \right\|_{L^2(4Q)}^p |Q|^{n(\frac{1}{p}-\frac{1}{2})p} + \sum_{j=3}^\infty \left\| S_1 \sqrt{L_1} b \right\|_{L^2(S_j(Q))}^p |2^j l(Q)|^{n(\frac{1}{p}-\frac{1}{2})p} \end{aligned}$$

$$\begin{aligned}
& \lesssim \left\| S_1 \sqrt{L_1} b \right\|_{L^2(4Q)}^p |Q|^{(\frac{1}{p}-\frac{1}{2})p} \\
& \quad + \sum_{j=3}^{\infty} \left\{ \iint_{\mathcal{R}(S_j(Q))} \left| t^{2k} L_1 e^{-t^{2k} L_1} b(y) \right|^2 \frac{dy dt}{t^{2k+1}} \right\}^{p/2} |2^j l(Q)|^{n(\frac{1}{p}-\frac{1}{2})p} \\
& \lesssim \left\| S_1 \sqrt{L_1} b \right\|_{L^2(4Q)}^p |Q|^{(\frac{1}{p}-\frac{1}{2})p} \\
& \quad + \sum_{j=3}^{\infty} \left\{ \int_{2^{j-2}Q} \int_{2^{j-3}l(Q)}^{\infty} \left| t^{2k} L_1 e^{-t^{2k} L_1} b(y) \right|^2 \frac{dy dt}{t^{2k+1}} \right\}^{p/2} |2^j l(Q)|^{n(\frac{1}{p}-\frac{1}{2})p} \\
& \quad + \sum_{j=3}^{\infty} \left\{ \int_{\mathbb{R}^n \setminus 2^{j-2}Q} \int_0^{\infty} \dots \right\}^{p/2} |2^j l(Q)|^{n(\frac{1}{p}-\frac{1}{2})p} \\
& =: \mathbf{I} + \sum_{j=3}^{\infty} (\mathbf{J}_j)^p + \sum_{j=3}^{\infty} (\mathbf{V}_j)^p.
\end{aligned}$$

For \mathbf{I} , by the $L^2(\mathbb{R}^n)$ -boundedness of S_1 , (6.10) and [6, Theorem 1.1], we see that

$$\mathbf{I} \lesssim \left\| \sqrt{L_1} b \right\|_{L^2(\mathbb{R}^n)}^p |Q|^{(\frac{1}{p}-\frac{1}{2})p} \lesssim \|b\|_{\dot{F}_{2,2}^k(\mathbb{R}^n)}^p |Q|^{(\frac{1}{p}-\frac{1}{2})p} \lesssim 1. \quad (6.13)$$

To estimate \mathbf{J}_j , recall the following *embedding theorem* (see, for example, [51]) that for all $f \in \dot{F}_{\frac{nr}{n+kr}, 2}^k(\mathbb{R}^n)$,

$$\|f\|_{L^r(\mathbb{R}^n)} \lesssim \|\nabla^k f\|_{L^{\frac{nr}{n+kr}}(\mathbb{R}^n)}. \quad (6.14)$$

For each \mathbf{O}_j , from Minkowski's inequality, Lemma 6.5, (6.14), Lemma 6.5, Hölder's inequality and (6.10), it follows that

$$\begin{aligned}
\mathbf{J}_j & \lesssim \left\{ \int_{2^{j-3}l(Q)}^{\infty} \left\| t^{2k} L_1 e^{-t^{2k} L_1} b \right\|_{L^2(2^{j-2}Q)}^2 \frac{dt}{t^{1+2k}} \right\}^{1/2} |2^j l(Q)|^{n(\frac{1}{p}-\frac{1}{2})} \\
& \lesssim \left\{ \int_{2^{j-3}l(Q)}^{\infty} t^{2n(\frac{1}{2}-\frac{1}{r})} \frac{dt}{t^{1+2k}} \right\}^{1/2} |2^j l(Q)|^{n(\frac{1}{p}-\frac{1}{2})} \|b\|_{L^r(Q)} \\
& \lesssim [2^j l(Q)]^{n(\frac{1}{p}-\frac{1}{r})-k} \|b\|_{L^r(Q)} \lesssim [2^j l(Q)]^{n(\frac{1}{p}-\frac{1}{r})-k} \|\nabla^k b\|_{L^{\frac{rn}{n+kr}}(Q)} \\
& \lesssim [2^j l(Q)]^{n(\frac{1}{p}-\frac{1}{r})-k} \|\nabla^k b\|_{L^2(Q)} |l(Q)|^{\frac{n+kr}{r}-\frac{n}{2}} \lesssim 2^{[n(\frac{1}{p}-\frac{1}{r})-k]j}.
\end{aligned}$$

Let $\alpha := \frac{n}{r} + k - \frac{n}{p}$. Since $p \in (\frac{nr}{n+kr}, 1]$, we then have $\alpha > 0$ and

$$\sum_{j=3}^{\infty} (\mathbf{J}_j)^p \lesssim \sum_{j=3}^{\infty} 2^{-\alpha j p} \lesssim 1. \quad (6.15)$$

To estimate \mathbf{V}_j , we write

$$\begin{aligned}
\mathbf{V}_j & \lesssim [2^j l(Q)]^{n(\frac{1}{p}-\frac{1}{2})} \left\{ \int_{\mathbb{R}^n \setminus 2^{j-2}Q} \int_0^{2^{j-3}l(Q)} \left| t^{2k} L_1 e^{-t^{2k} L_1} b(y) \right|^2 \frac{dy dt}{t^{2k+1}} \right\}^{1/2} \\
& \quad + [2^j l(Q)]^{n(\frac{1}{p}-\frac{1}{2})} \left\{ \int_{\mathbb{R}^n \setminus 2^{j-2}Q} \int_{2^{j-3}l(Q)}^{\infty} \dots \right\}^{1/2} \\
& =: \mathbf{V}_{j,1} + \mathbf{V}_{j,2}.
\end{aligned}$$

Similar to the estimate of J_j , we see that

$$V_{j,2} \lesssim 2^{-\alpha j}. \quad (6.16)$$

To estimate $V_{j,1}$, let $\beta \in (2k + 2n(1/r - 1/2), \infty)$. By Lemma 6.5, (6.10) and (6.14), there exists a positive constant \tilde{C} such that

$$\begin{aligned} V_{j,1} &\lesssim [2^j l(Q)]^{n(\frac{1}{p}-\frac{1}{2})} \left[\int_0^{2^{j-3}l(Q)} t^{2n(\frac{1}{2}-\frac{1}{r})} \exp \left\{ -\tilde{C} \frac{[2^j l(Q)]^{2k/(2k-1)}}{t^{2k/(2k-1)}} \right\} \frac{dt}{t^{2k+1}} \right]^{1/2} \|b\|_{L^r(Q)}, \\ &\lesssim [2^j l(Q)]^{n(\frac{1}{p}-\frac{1}{2})} \left[\int_0^{2^{j-3}l(Q)} t^{2n(\frac{1}{2}-\frac{1}{r})} \left[\frac{t}{2^j l(Q)} \right]^\beta \frac{dt}{t^{2k+1}} \right]^{1/2} \|\nabla^k b\|_{L^{\frac{rn}{n+kr}}(Q)} \\ &\lesssim [2^j l(Q)]^{n(\frac{1}{p}-\frac{1}{2})} \left[\frac{1}{[2^j l(Q)]^\beta} \int_0^{2^{j-3}l(Q)} t^{2n(\frac{1}{2}-\frac{1}{r})+\beta-2k-1} dt \right]^{1/2} \|\nabla^k b\|_{L^2(Q)} |l(Q)|^{\frac{n+kr}{r}-\frac{n}{2}} \\ &\lesssim 2^{j[n(\frac{1}{p}-\frac{1}{r})-k]}, \end{aligned}$$

which, together with (6.16), shows that

$$\sum_{j=3}^{\infty} (V_j)^p = \sum_{j=3}^{\infty} 2^{-\alpha j p} \lesssim 1.$$

This, combined (6.13) and (6.15), implies (6.12), which completes the proof of Proposition 6.6. \square

Combining Theorem 6.2 and Proposition 6.6, we obtain the following Riesz transform characterization of $H_{L_1}^p(\mathbb{R}^n)$. We point out that Theorem 6.10 when $k = 1$ is just the Riesz transform characterization of $H_{-\text{div}(A\nabla)}^p(\mathbb{R}^n)$ for $p \in (0, 1]$, which is exactly [39, Theorem 5.2] in the case that $p \in (0, 1]$.

Theorem 6.10. *Let $k \in \mathbb{N}$, L_1 be the $2k$ order divergence form homogeneous elliptic operator, $r \in (1, 2]$ such that $rn/(n+kr) \leq 1$, and the semigroup $\{e^{-tL_1}\}_{t>0}$ satisfy the $L^r - L^2$ k -off-diagonal estimates. Then for all $p \in (rn/(n+kr), 1]$, $H_{L_1}^p(\mathbb{R}^n) = H_{\text{Riesz}, L_1}^p(\mathbb{R}^n)$ with equivalent norms.*

Remark 6.11. We point out that a key fact used in the proof of Proposition 6.6 (and hence Theorem 6.10) is $\|\sqrt{L_1}f\|_{L^2(\mathbb{R}^n)} \lesssim \|\nabla^k f\|_{L^2(\mathbb{R}^n)}$, which comes from [6, Theorem 1.1]. This inequality for L_2 is equivalent to the following inequality that for all $f \in \dot{W}^{k,2}(\mathbb{R}^n)$,

$$\left\| V^{k/2} f \right\|_{L^2(\mathbb{R}^n)} \lesssim \left\| \nabla^k f \right\|_{L^2(\mathbb{R}^n)},$$

which seems impossible even when $V := 1$. Thus, the method used in the proof of Proposition 6.6 seems unsuitable for obtaining a counterpart of Proposition 6.6 for L_2 .

Acknowledgements This work was supported by the National Natural Science Foundation (Grant No. 11171027) of China and Program for Changjiang Scholars and Innovative Research Team in University of China. The authors would like to thank the referees for their careful reading and many valuable remarks which made this article more readable.

References

- 1 Ahn B, Li J. Orlicz-Hardy spaces associated to operators satisfying bounded H_∞ functional calculus and Davies-Gaffney estimates. J Math Anal Appl, 2011, 373: 485-501
- 2 Albrecht D, Duong X T, McIntosh A. Operator theory and harmonic analysis. In: Instructional Workshop on Analysis and Geometry, Part III (Canberra, 1995), Proc Centre Math Appl Austral Nat Univ, vol. 34. Canberra: Austral Nat Univ, 1996, 77-136

- 3 Auscher P. On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates. *Mem Amer Math Soc*, 2007, 871: 1-75
- 4 Auscher P, Duong X T, McIntosh A. Boundedness of Banach space valued singular integral operators and Hardy spaces. Unpublished preprint, 2005
- 5 Auscher P, Hofmann S, Lacey M, et al. The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n . *Ann of Math* (2), 2002, 156: 633-654
- 6 Auscher P, Hofmann S, McIntosh A, et al. The Kato square root problem for higher order elliptic operators and systems on \mathbb{R}^n . *J Evol Equ*, 2001, 1: 361-385
- 7 Auscher P, McIntosh A, Russ E. Hardy spaces of differential forms on Riemannian manifolds. *J Geom Anal*, 2008, 18: 192-248
- 8 Auscher P, Russ E. Hardy spaces and divergence operators on strongly Lipschitz domains of \mathbb{R}^n . *J Funct Anal*, 2003, 201: 148-184
- 9 Barbatis G, Davies E. Sharp bounds on heat kernels of higher order uniformly elliptic operators. *J Operator Theory*, 1996, 36: 179-198
- 10 Blunck S, Kunstmann P C. Weak type (p, p) estimates for Riesz transforms. *Math Z*, 2004, 247: 137-148
- 11 Blunck S, Kunstmann P C. Generalized Gaussian estimates and the Legendre transform. *J Operator Theory*, 2005, 53: 351-365
- 12 Cao J, Liu Y, Yang D. Hardy spaces $H^1_L(\mathbb{R}^n)$ associated to Schrödinger type operators $(-\Delta)^2 + V^2$. *Houston J Math*, 2010, 36: 1067-1095
- 13 Chang D -C, Dafni G, Stein E M. Hardy spaces, BMO and boundary value problems for the Laplacian on a smooth domain in \mathbb{R}^n . *Trans Amer Math Soc*, 1999, 351: 1605-1661
- 14 Chang D -C, Krantz S G, Stein E M. Hardy spaces and elliptic boundary value problems. *Contemp Math*, 1992, 137: 119-131
- 15 Chang D-C, Krantz S G, Stein E M. H^p theory on a smooth domain in \mathbb{R}^N and elliptic boundary value problems. *J Funct Anal*, 1993, 114: 286-347
- 16 Cho Y, Kim J. Atomic decomposition on Hardy-Sobolev spaces. *Studia Math*, 2006, 177: 25-42
- 17 Coifman R R, Meyer Y, Stein E M. Some new function spaces and their applications to harmonic analysis. *J Funct Anal*, 1985, 62: 304-335
- 18 Coifman R R, Weiss G. Extensions of Hardy spaces and their use in analysis. *Bull Amer Math Soc*, 1977, 83: 569-645
- 19 Davies E B. Uniformly elliptic operators with measurable coefficients. *J Funct Anal*, 1995, 132: 141-169
- 20 Duong X T, Li J. Hardy spaces associated to operators satisfying bounded H_∞ functional calculus and Davies-Gaffney estimates. Preprint
- 21 Duong X T, Xiao J and Yan L. Old and new Morrey spaces with heat kernel bounds. *J Fourier Anal Appl*, 2007, 13: 87-111
- 22 Duong X T, Yan L. New function spaces of BMO type, the John-Nirenberg inequality, interpolation, and applications. *Comm Pure Appl Math*, 2005, 58: 1375-1420
- 23 Duong X T, Yan L. Duality of Hardy and BMO spaces associated with operators with heat kernel bounds. *J Amer Math Soc*, 2005, 18: 943-973
- 24 Dziubański J, Zienkiewicz J. Hardy space H^1 associated to Schrödinger operator with potential satisfying reverse Hölder inequality. *Rev Mat Iberoamericana* 1999, 15: 279-296
- 25 Dziubański J, Zienkiewicz J. H^p spaces for Schrödinger operators. In: *Fourier analysis and related topics* (Bpolhk edlewo, 2000), Banach Center Publ, vol. 56. Warsaw: Polish Acad Sci, 2002, 45-53
- 26 Fefferman C, Stein E M. H^p spaces of several variables. *Acta Math*, 1972, 129: 137-193
- 27 García-Cuerva J, Rubio de Francia J. *Weighted Norm Inequalities and Related Topics*. Amsterdam: North-Holland Publishing Co, 1985
- 28 Grafakos L. *Classical Fourier Analysis*. New York: Springer Press, 2008
- 29 Grafakos L, Liu L, Yang D. Boundedness of paraproduct operators on RD-spaces. *Sci China Math*, 2010, 53: 2097-2114
- 30 Grafakos L, Liu L, Yang D. Maximal function characterizations of Hardy spaces on RD-spaces and their applications. *Sci China Ser A*, 2008, 51: 2253-2284
- 31 Li B, Bownik M, Yang D, et al. Anisotropic singular integrals in product spaces. *Sci China Math*, 2010, 53: 3163-3178
- 32 Liang Y, Yang D, Yang S. Applications of Orlicz-Hardy spaces associated with operators satisfying Poisson estimates. *Sci China Math*, 2011, 54: 2395-2426
- 33 Haase M. *The Functional Calculus for Sectorial Operators*. In: *Oper Theory Adv Appl*, vol. 169. Basel: Birkhäuser Verlag, 2006
- 34 Han Y, Paluszynski M, Weiss G. A new atomic decomposition for the Triebel-Lizorkin spaces. In: *Harmonic analysis and operator theory* (Caracas, 1994), *Contemp Math*, vol. 189. Providence, RI: Amer Math Soc, 1995, 235-249

- 35 Hofmann S, Lu G, Mitrea D, et al. Hardy spaces associated to non-negative self-adjoint operators satisfying Davies-Gaffney estimates. *Mem Amer Math Soc*, 2011, 1007: 1-78
- 36 Hofmann S, Martell J. L^p bounds for Riesz transforms and square roots associated to second order elliptic operators. *Publ Mat*, 2003, 47: 497-515
- 37 Hofmann S, Mayboroda S. Hardy and BMO spaces associated to divergence form elliptic operators. *Math Ann*, 2009, 344: 37-116
- 38 Hofmann S, Mayboroda S. Correction to "Hardy and BMO spaces associated to divergence form elliptic operators". arXiv: 0907.0129
- 39 Hofmann S, Mayboroda S, McIntosh A. Second order elliptic operators with complex bounded measurable coefficients in L^p , Sobolev and Hardy spaces. *Ann Sci École Norm Sup (4)*, 2011, 44: 723-800
- 40 Hu G, Yang D, Zhou Y. Boundedness of singular integrals in Hardy spaces on spaces of homogeneous type. *Taiwanese J Math*, 2009, 13: 91-135
- 41 Jiang R, Yang D. Orlicz-Hardy spaces associated with operators satisfying Davies-Gaffney estimates. *Commun Contemp Math*, 2011, 13: 331-373
- 42 Jiang R, Yang D. Predual spaces of Banach completions of Orlicz-Hardy spaces associated with operators. *J Fourier Anal Appl*, 2011, 17: 1-35
- 43 Jiang R, Yang D. New Orlicz-Hardy spaces associated with divergence form elliptic operators. *J Funct Anal*, 2010, 258: 1167-1224
- 44 Jiang R, Yang D. Generalized vanishing mean oscillation spaces associated with divergence form elliptic operators. *Integral Equations Operator Theory*, 2010, 67: 123-149
- 45 Jiang R, Yang D, Zhou Y. Orlicz-Hardy spaces associated with operators. *Sci China Ser A*, 2009, 52: 1042-1080
- 46 McIntosh A. Operators which have an H_∞ functional calculus. In: *Miniconference on operator theory and partial differential equations (North Ryde, 1986)*, *Proc Centre Math Anal Austral Nat Univ*, vol. 14. Canberra: Austral Nat Univ, 1986, 210-231
- 47 Ouhabaz E M. Analysis of Heat Equations on Domains. In: *London Mathematical Society Monographs Series 31*. Princeton, NJ: Princeton University Press, 2005
- 48 Stein E M. Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals. Princeton, NJ: Princeton University Press, 1993
- 49 Stein E M, Weiss G. On the theory of harmonic functions of several variables. I. The theory of H^p -spaces. *Acta Math*, 1960, 103: 25-62
- 50 Taibleson M H, Weiss G. The molecular characterization of certain Hardy spaces. In: *Astérisque*, 77, Representation theorems for Hardy spaces. Paris: Soc Math France, 1980, 67-149
- 51 Triebel H. Theory of Function Spaces. Basel: Birkhäuser Verlag, 1983
- 52 Welland G, Zhao S. ϵ -families of operators in Triebel-Lizorkin and tent spaces. *Canad J Math*, 1995, 47: 1095-1120
- 53 Yan L. Classes of Hardy spaces associated with operators, duality theorem and applications. *Trans Amer Math Soc*, 2008, 360: 4383-4408
- 54 Yang Da, Yang Do. Boundedness of linear operators via atoms on Hardy spaces with non-doubling measures. *Georgian Math J*, 2011, 18: 377-397
- 55 Yang D C, Yang S B. Local Hardy spaces of Musielak-Orlicz type and their applications. *Sci China Math*, 2012, 55, doi: 10.1007/s11425-012-4377-z (in press) or arXiv:1108.2797
- 56 Yang D, Zhou Y. Localized Hardy spaces H^1 related to admissible functions on RD-spaces and applications to Schrödinger operators. *Trans Amer Math Soc*, 2011, 363: 1197-1239